

**Surveillance of Viral Hepatitis in Hong Kong**  
**- 2017 Update Report**

**Viral Hepatitis Control Office  
Special Preventive Programme  
Centre for Health Protection  
Department of Health  
December 2018**

*The information contained in this Report is up to year 2017 for the surveillance data, service statistics and published research findings.*

### **Editorial Team:**

Dr Kenny Chan  
Dr Rebecca Lam  
Dr Bonnie Wong  
Dr Priscilla Kwok  
Ms Wai Kit Chan  
Ms Phoebe Lam

### **Correspondence**

Viral Hepatitis Control Office  
Special Preventive Programme  
9/F, Kowloon Bay Health Centre  
9 Kai Yan Street  
Kowloon  
Hong Kong

Telephone: (852) 2116 2888

Facsimile: (852) 2117 0809

Website: [www.hepatitis.gov.hk](http://www.hepatitis.gov.hk)

E-mail: [hepatitis@dh.gov.hk](mailto:hepatitis@dh.gov.hk)

Comments and suggestions on this Report are most welcome.

\*pdf version of the report can be downloaded from [www.hepatitis.gov.hk](http://www.hepatitis.gov.hk).

# SURVEILLANCE OF VIRAL HEPATITIS IN HONG KONG

2017 UPDATE

## CONTENTS

<b>CONTENTS</b>	<b>3</b>
<b>ACKNOWLEDGEMENTS</b>	<b>4</b>
<b>COMMENTARY</b>	<b>5</b>
<b>TABULATED RESULTS</b>	
<b>Acute viral hepatitis under the disease notification system</b>	<b>23</b>
<b>Seroprevalence of hepatitis A and hepatitis E</b>	<b>42</b>
<b>Seroprevalence and vaccination coverage of hepatitis B</b>	<b>48</b>
<b>Seroprevalence of hepatitis C</b>	<b>72</b>
<b>Liver cancers</b>	<b>80</b>
<b>ABBREVIATIONS</b>	<b>85</b>
<b>REFERENCES</b>	<b>86</b>

# SURVEILLANCE OF VIRAL HEPATITIS IN HONG KONG

## 2017 UPDATE

### ACKNOWLEDGEMENTS

The Special Preventive Programme wishes to thank the following agencies for their contributions to the preparation of the 2017 Update Report:

Action for REACH OUT

CHC-Group Medical Practice

Chinese University of Hong Kong

Department of Medicine, Princess Margaret Hospital

Department of Microbiology, Prince of Wales Hospital

Department of Microbiology, Princess Margaret Hospital

Family Health Service, Department of Health

Family Planning Association of Hong Kong

Public Health Laboratory Services Branch, Centre for Health Protection, Department of Health

Health Service of Baptist University of Hong Kong

Hong Kong Red Cross Blood Transfusion Service

Hong Kong Cancer Registry

Pamela Youde Nethersole Eastern Hospital

Surveillance & Epidemiology Branch, Centre for Health Protection, Department of Health

TB and Chest Service, Centre for Health Protection, Department of Health

The University of Hong Kong

# SURVEILLANCE OF VIRAL HEPATITIS IN HONG KONG

## 2017 UPDATE

### COMMENTARY

#### Surveillance Mechanisms of Viral Hepatitis in Hong Kong

1. Viral hepatitis is a statutory notifiable disease in Hong Kong. Locally, voluntary reporting was started in as early as 1966 and, since 1974, the disease has become notifiable. It was not until 1988 that the reported cases were classified by viral etiology, namely hepatitis A, hepatitis B, non-A non-B hepatitis and unclassified hepatitis. In 1996, non-A non-B hepatitis was further categorised into hepatitis C, hepatitis E and hepatitis (not elsewhere classified).

2. The extent of chronic infections resulting from some hepatitis, notably hepatitis B and C, is determined by other mechanisms. Insight into the epidemiology of various forms of hepatitis in Hong Kong can be gained by analytical interpretation of regular statistics collected by health care or other institutions, and information generated from various studies. This Report presents the latest findings from collation and analysis of viral hepatitis data obtained from the disease notification system, service statistics, seroprevalence studies and other research findings. Much as one hopes that the local viral hepatitis picture can be painted accurately and fully, this is certainly limited by the nature and availability of data. The presence of biases in data *per se* and their interpretation need to be acknowledged in reading this Report.

#### Epidemiology of HAV and HEV

3. Hepatitis A virus (HAV) and hepatitis E virus (HEV) are both transmitted by the faecal-oral route, albeit with different local epidemiology in the past two decades. Hong Kong was once of intermediate endemicity for HAV [1, 2]. After 1988 when viral hepatitis began to be reported according to etiologic agents, the largest epidemic of hepatitis A occurred in 1992, with over 3,500 cases reported to the Department of Health (DH) (Box 1). This represented a notification rate of 63 per 100,000 population (Box 7) and since then, a gradual declining trend in HAV incidence has been observed. This discernible decline in hepatitis A contributed to a parallel declining trend in overall

reported viral hepatitis since 2002 (Box 3). The case fatality rates from hepatitis A has been low and ranged between 0 and 0.15% in the last two decades (Box 7).

4. From 2008 to 2017, the annual number of hepatitis A cases reported ranged from 43 to 138 (Box 1). In 2015, a review on 587 reported cases of hepatitis A from 2005 to 2014 was published by the Surveillance and Epidemiology Branch (SEB) of Centre for Health Protection (CHP), Department of Health (DH). The male to female ratio was 1.2 to 1, with 75% aged below 40 years. The majority (70%) of cases required hospitalisation, and two fatal cases were recorded. Both fatalities had multiple comorbidities. The majority (76%) of the patients acquired the disease locally. 92% were sporadic cases and 22 small clusters affecting two to four patients were identified, at least 60% of which were clusters affecting members of the same household [3].

5. An increase in the number of cases was noted in 2015 when a total of 138 cases were reported. The majority (75%) of the cases was reported from February to June. The male to female ratio was 1.2 to 1, with a median age of 33 years (range: 3 to 83 years). There was no fatality. Except two cases studying in the same school and two cases from same family, no epidemiological link was found. No single identifiable source could explain the upsurge of cases [3].

6. In 2016, a total of 98 cases of hepatitis A were recorded, affecting 68 men and 30 women (male to female ratio 2.26:1) aged from 3 to 86 years (median 32 years). Sixty-three cases (64.3%) acquired the infection locally. 85.7% required hospitalisation.

7. In late 2016, an unusual upsurge of acute hepatitis A infection affecting men who have sex with men (MSM) with human immunodeficiency virus infection was noticed. With retrospective investigations and prospective reporting, a total of 53 cases of laboratory-confirmed HAV infection with clinical symptoms among individuals identified as MSM were recorded between September 2015 and November 2017. The age range was 20 to 55 years (median: 33 years). Forty-five (84.9%) required hospitalisation and there were no fatalities. Thirty-seven cases (69.8%) were known to be HIV-positive attending one of the three designated public HIV clinics. The majority (96.2%) did not report history of HAV vaccination. Eighteen (33.9%) reported travel history within the incubation period. Around one quarter of the cases had concurrent diagnosis of other sexually transmitted infections including syphilis, gonorrhoea and chlamydia infection. Among the cases with specimen available for laboratory analysis, forty-three (81.1%) had identical nucleotide sequences within the

genotyping window. Apart from one cluster affecting two patients who were sex partners residing together, no other epidemiological linkage could be found. No common food nor water source or social gathering was identified among these cases. Epidemiological investigations suggested that the outbreak was contributed by transmission by way of sexual contact between men, a high proportion of whom were HIV-infected. HAV outbreaks among MSM communities were being reported during the same period in some other regions with low HAV endemicity including Taiwan, Europe and both North and South America [4].

8. Over the years, there has been an increase in the proportion of reported cases over 35 years old. Although the majority were still below 44 years of age, the proportion of reported cases that were aged 45 and above had increased from less than 10% in the last two decades to 14%-25% since 2010 (Box 6).

9. In a local household study conducted in 2001, (Community Research Project for Viral Hepatitis 2001, CRPVH), anti-HAV positivity was less frequent ( $P < 0.001$ ) across all age groups among subjects  $> 21$  years old [2] than subjects in the same age groups of another study conducted in late 1980s [5]. HAV prevalence has only increased insignificantly in every 10-year age groups of people aged 21-50 when compared with their corresponding 10-year younger age groups, signifying an aging cohort effect with no major infections in the prior 10 years [2]. Similar conclusions can be drawn when comparing the late 1980s findings with those of a late 1970s study on local HAV seroprevalence [6]. Together, these three studies suggest that age-specific prevalence of HAV has right-shifted locally since 1980s. As of 2001, anti-HAV was present in about 20% of adults below 30 years old while it was over 80% in people aged  $\geq 40$  years in the general Chinese population (Box 20). Data from laboratory surveillance performed by Public Health Laboratory Services Branch (PHLSB) every five years had also shown that the seroprevalence of anti-HAV had remained below 40% among those younger than 30 years old in 2000, 2005 and 2010. From the available data, the prevalence of hepatitis A infection has been falling in Hong Kong, which echoes the finding of a higher median age in reported HAV cases that also reflects the increased susceptibility of the adult population. The data also indicates that Hong Kong has changed from a region with intermediate to very low endemicity in the past three decades (Box 21) [7].

10. Besides an increasing prevalence with higher age, people born outside Hong Kong were generally more likely to test positive for anti-HAV whereas the reverse was true for people of non-labour work [2]. From the telephone interview part of the CRPVH 2001, some 11% of 4,564 subjects reported a history of HAV vaccination, about 80% of whom had completed the course. More people less than 40 years old had received the vaccination. Over 98% had the cost paid by themselves or covered by their employers. In the latest serosurvey conducted by PHLSB in 2015, there was a significant increase in the seroprevalence of anti-HAV in the younger age group, most prominent among those aged 0-10 (Box 21). These findings may suggest an increase in uptake of HAV vaccination in the community.

11. Cross-sectional surveys of anti-HAV at Kowloon Bay Integrated Treatment Centre (ITC), the HIV specialist clinic under Department of Health, have been started since 2007. The subjects consisted of all new HIV/AIDS patients who first attended ITC between Jul 2007 and 2017 and convenience samples of all active HIV/AIDS patients who first attended ITC before Jul 2007 (Box 22). The prevalence of anti-HAV increased with age of HIV/AIDS patients, and the overall positivity rate among these patients tested between 2007 and 2017 appeared to be comparable with that of the 2010 serosurvey data. Confounding factors, such as different levels of past infection, immunodeficiency in HIV patients, history of HAV vaccination and difference in years of testing, may have affected the results. Compared with patients acquiring HIV via other routes, those infected via homosexual or bisexual routes were at the highest risk of hepatitis A infection, as reflected by the lowest level of anti-HAV prevalence in this group of patients (Box 23). Indeed, the increased susceptibility had manifested itself during the upsurge of hepatitis A infection among MSM occurring in 2015 to 2017 [4]. As a result, the Scientific Committee on AIDS and STI, and Scientific Committee on Vaccine Preventable Diseases had extended their recommendation for hepatitis A vaccine to MSM in June 2017 [8].

12. The annual notification of hepatitis E infection increased from 11 in 1996 to a record high of 150 in 2012 (Box 1). In the past five years, the number of reported cases of hepatitis E ranged from 64 to 96. A seasonal pattern was observed with peak infections reported from February to April (Box 15), indicating that infection was more common during winter and spring seasons. Of 1271 cases reported, 838 (65.9%, Box 16) were male, giving male to female ratio of 1.9:1. The majority was adults, most of whom were aged between 35 to 74 (Box 17). Fatalities were more common with acute



hepatitis E than with acute hepatitis A and the death rate reached as high as 0.44 per million population in 2002 when three deaths attributable to acute hepatitis E infection had occurred (Box 18).

13. In the CRPVH study conducted in 2001, 18.8% of adult subjects were found to have serologic evidence of HEV infection. People in the 40-49 years age group had the highest positivity rate of 24.1% (Box 24). Another local seroprevalence study on anti-HEV using 450 serum samples submitted for virological investigation in 2008-2009 in a local hospital found a higher rate of HEV IgG seropositivity [9]. The HEV IgG seropositivity rate increased from 8% among 1-10 years old to >56% among those aged over 80. The overall seropositivity rate was higher among male than female (32.9% vs 24.4%,  $p=0.048$ ). Despite the limitations of small sample size and bias sampling in this study, the finding of an overall increase in the seropositivity rate is compatible with the changing local epidemiology of Hepatitis E notified to Department of Health in recent years.

14. The CHP reviewed all hepatitis E cases recorded between 2001 to 2010 [10]. Of the 524 cases, the commonest presentations were tea-coloured urine, jaundice, anorexia, fever, myalgia and nausea. 78.2% were hospitalised with a median stay of 7 days. A total of 12 cases were fatal (9 males and 3 females), age ranged from 53 to 82 (median age 67.5 years). The case fatality rate was 2.3%, which was comparable with reported figures from other countries. None of the fatal cases were pregnant. Most cases (99.4%) were sporadic infection and 87.4% acquired the disease locally. A small family cluster involving 2 males (aged 15 and 44 years) was identified. The 2 victims had shared multiple high-risk food items at home during the incubation period. It proved difficult to determine the exact source of infection of individual sporadic cases as hepatitis E has a long incubation period of 15-64 days. Nonetheless, epidemiological investigation has not identified any outbreak linked to a particular food premises.

15. Another review of the acute hepatitis E cases recorded by CHP from 2012 to May 31 2017 showed a total of 554 cases, with age ranging from 15 to 96 years (median: 55 years) [11]. More males were affected than females (63.0% vs. 37.0%). More cases were recorded in February and March. Most of the cases (83.8%) acquired the infection locally. Symptomatology was similar with the cases from 2001 to 2010. Four hundred and seventy-six (85.9%) patients required hospitalization with a median length of stay of six days. Nine fatal cases were recorded, among whom 7 had

underlying illnesses, giving a case fatality rate of 1.6%. The age of the deceased patients ranged from 59 to 79 years (median: 74 years). All cases were sporadic infections, except for a pair of father and son who shared meals frequently during the incubation period. None of the cases was related to outbreak involving food premises. A significant proportion of the victims recalled consuming pig offals (37.5%) and shellfish (29.4%) during the incubation period.

16. Another published study identified differences in epidemiology and clinical features between sporadic hepatitis E and hepatitis A cases. Of 105 acute hepatitis A and 24 hepatitis E patients seen at Princess Margaret Hospital (PMH) in 2002, HAV patients were significantly younger (median age of 27 years) and had recent history of shellfish consumption while HEV patients were older (median age = 53 year) and most had a recent travel history. Moreover, whereas hepatitis A was milder and recovery was uneventful, hepatitis E was more severe, associated with significant mortality and frequently complicated by protracted coagulopathy and cholestasis [12].

17. A local study examined the genotype of 57 patients with acute HEV infection who were admitted to Prince of Wales Hospital (PWH). Fifty-six patients (98%) were Chinese. All cases were sporadic. No fulminant hepatitis was recorded and all patients recovered. Phylogenetic analyses of the open reading frame ORF2 fragments from 46 patients and ORF1 fragments from 33 patients showed complete agreement, with most (n= 45 [98%]) belonging to genotype 4. The remaining isolate was genotype 3 obtained from a woman who had no history of travel. Most of the Hong Kong isolates clustered closely with a swine isolate reported from Guangxi Province, China [13].

18. Apart from pregnancy, coinfection with chronic Hepatitis B virus might be associated with more fulminant clinical outcome in patients infected with Hepatitis E. Among 3 cases of serious infection of Hepatitis E with acute liver failure reported to DH in the first two months of 2012, one required liver transplantation and two passed away. One of the deceased patients was tested positive for chronic hepatitis B infection [14]. Moreover, a 10-year retrospective study on acute hepatitis E in local hospitals showed that patients with chronic HBV acutely infected with hepatitis E had a higher rate of liver failure, liver-related mortality and all-cause mortality, though the association was not statistically significant [15].

19. There is evidence suggesting a zoonotic source of Hepatitis E in overseas studies, and that pigs may be an important reservoir. In light of these observations, the Centre for Food Safety conducted a risk assessment study titled "Hepatitis E Virus in Fresh

Pig Livers” [16] to determine the HEV prevalence in fresh pig liver samples obtained in local markets. One hundred fresh pig liver samples were collected from pigs slaughtered between mid-January and May. Sixteen (31%) out of 51 roaster pig (around four months old) liver samples were positive for HEV, while none of the 49 porker pig (around six months old) liver samples tested positive. Partial sequences of some HEV isolates from roaster pigs were identical to those from 7 among 48 local human cases with date of onset from January to July 2009, as well as local cases recorded in the past. The findings suggest the possibility of roaster pigs as one of the sources of local human hepatitis E infections.

20. One HEV vaccine was licensed in China in December 2011 for use in people aged at or older than 16 years old [17]. To date, it has not been licensed in other countries or territories. It has been shown to have high efficacy against hepatitis E in healthy adults of 16 to 65 years old in China. Data is however limited on its impact on the overall disease incidence and reduction of mortality in the general population where disease is endemic. Therefore, in the absence of sufficient information, World Health Organization (WHO) has not made recommendation on its incorporation in national programmes [17].

### **Pattern of Hepatitis B in Various Communities and its Significance**

21. The number of reported acute hepatitis B virus (HBV) infections has been decreasing over the last decade, from 137 cases reported in 2000 to 33 cases reported in 2017 (Box 1). In an epidemiologic study of acute HBV by the Department of Health and Hong Kong Red Cross Blood Transfusion Service (HKRCBTS), 149 of 351 eligible subjects recruited from 2000 to 2003 participated in risk factor assessment with or without blood screening. Repeat blood donors who tested positive for HBsAg for the first time and were then confirmed IgM anti-HBc positive were reported as having acute HBV. There were 43 such clients, yielding an incidence rate of HBV seroconversion in repeat donors as 9.4/100,000 (n=148,366), 9.3/100,000 (n=150,420), 4.6/100,000 (n=151,410) and 3.5/100,000 (n=143,230) in 2000, 2001, 2002 and 2003 respectively. Nearly 70% of the study subjects were male; 99% were Chinese and the mean age was 31 years. Over half could not have risk factor of acute HBV determined despite undergoing a standardised questionnaire interview by nurses. Sexual contact was assessed to be the commonest risk (85%) in the rest. Of 124 subjects who had hepatitis B screening at 6 months post-IgM anti-HBc positivity, 50% developed anti-HBs while 9.7% were HBsAg positive. Although these results could suggest a higher

rate of HBV chronicity than what was previously reported in the literature, they have to be interpreted with caution owing to the relative small number of samples, incompleteness of data and potential biases from the subjects sampling and other study design.

22. Determining the seroprevalence of hepatitis B surface antigen (HBsAg) sheds light on how common chronic HBV infection is in different communities, as well as informing its chronic disease burden. The various adult communities can be categorised into 3 groups according to the risk of contracting HBV:

- (a) without apparent risk: blood donors, pre-marital/ pre-pregnancy service users, antenatal women, police officers, new health care workers (HCW)
- (b) with undetermined risk: clients seeking post-exposure management and tuberculosis patients
- (c) with apparent risk: drug users, HIV/AIDS patients and female sex workers

23. A majority of the available seroprevalence data in different populations were limited to overall positivity rate of HBV markers. Still, temporal trend can be discerned as most have yearly data for the past decade or so. Several features on the current pattern of HBV could be observed from the serologic investigations, namely

- (a) chronic HBV infection is in a general declining trend in community groups without apparent risk of contracting HBV
- (b) HBV prevalence increases with increasing age, and
- (c) chronic HBV infection is commoner in male than female.

24. A word of caution in the interpretation of data though, is that testing for HBV markers has been performed for a variety of reasons in different communities, with heterogeneous mix of population characteristics.

### **Seroprevalence of adult communities without apparent risks**

25. The temporal decline of chronic HBV infection has been most obvious in new blood donors and police officers. For new blood donors, the HBsAg prevalence follows a continual falling trend since early 1990s, from 8% in 1990 to 0.9% in year 2017 (Box 25). The trend is even more obvious among the 16-19 years age group where the prevalence is as low as 0.3% in both female and male (Box 26). A similar trend was observed among police officers where the HBsAg prevalence falls from 7.9% in 1997

to 1.4% in 2017 (Box 32), with the lowest prevalence of 0.5% among those aged 20 or less (Box 33). A falling trend was also observed in other community groups without apparent HBV risk, albeit less prominent (Box 42, 43).

26. The HBsAg prevalence in antenatal mothers has been decreasing from over 10% in the early 1990s to 5.0% in 2017 (Box 29). As compared with other groups without apparent risk, the overall HBsAg prevalence in antenatal mothers is higher and confounded by the place of birth. A study of 2480 pregnant women attending the Maternal and Child Health Centre (MCHC) of DH in 1996 found a 13.1% in those born in Mainland China as compared to 8.4% in local mothers [18]. Data from Virus Unit, Department of Health also showed a higher prevalence of 12.5% and 13.8% in the subset of non-resident expectant mothers versus the overall positivity rate of 8.5% and 8.6% in 2004 and 2005 respectively. The prevalence in pre-marital/ pre-pregnancy package service users has dropped from 9.6% in 1990 to remain static in the range of 4.8% to 6.9% in the past decade (Box 28). The prevalence in newly recruited health care workers as determined at pre-HBV vaccination screening also showed a decreasing trend from 5.9% in 2001 to 3.5% in 2017 among female, and from 6.1% in 2001 to 2.4% in 2017 among male (Box 35).

#### **Seroprevalence of adult communities with undetermined risk**

27. Of 874 tuberculosis patients attended TB & Chest Clinics, DH between March and May in 2015, 74 (8.5%, Box 36) were detected HBsAg positive, with the highest prevalence rate in the middle age group (40-59 years old: 10%, Box 37) followed by the more elderly group ( $\geq 60$  years old: 8.8%, Box 37). The HBsAg positivity rate was also found to be higher in male clients (10%) than in female (5.7%, Box 36). Both the age (Box 37) and gender pattern (Box 36) were consistently observed over the last decade. Among clients attended for post exposure management, HBsAg rate was found higher in non-health care workers than in health care workers (Box 38), which may be partly explained by the success of pre-employment vaccination programme for healthcare workers.

#### **Seroprevalence of adult communities with apparent risk**

28. The HBsAg prevalence in HIV/AIDS patients under care of DH was in the range of 5.6% to 11.3% in the past decade (Box 40). Due to underlying immunosuppression and shared routes of transmission, HIV/AIDS patients are more likely to be chronically infected with HBV [19]. The HBsAg prevalence in female sex workers attending the

clinic of Action for REACH OUT tested between 2007 to 2011 ranged from 5.0% to 10.4% (Box 43). The data regarding prevalence of HBsAg in drug users was difficult to interpret because of the small number of subjects since 2006 (Box 39). Overall, the difference in HBsAg prevalence between groups with or without apparent risk of contracting HBV has not been prominent in the past few years.

### Genotypes of Hepatitis B and their Disease Course

29. Different HBV genotypes have been identified with distinct geographic distribution and association with different clinical outcomes. Local studies indicated that genotype C was the commonest genotype and genotype B was the second. A study of 776 chronic hepatitis B patients seen at the University of Hong Kong Liver clinic from 1999 to mid-2003 found that genotype C was the commonest (486, 62.6%), followed by genotype B (252, 32.5%), with a majority of genotype B belonged to subgroup Ba [20]. Another study of 426 chronic HBV patients recruited consecutively from 1997 to mid-2000 at the Hepatitis clinic of Prince of Wales Hospital (PWH) found a prevalence of 57% (242) and 42% (179) of genotypes C and B respectively [21].

30. A study of 49 HBV genotype C ethnic Chinese patients under the care of the PWH Hepatitis Clinic identified 2 distinct groups with different epidemiological distribution and virologic characteristics – 80% being genotype “Cs” (found mostly in Southeast Asia) and 20% “Ce” (predominated in Far East) [22]. In addition, subgenotype Cs appears to be more common in Hong Kong than other parts of China. In the recent analysis of a cohort of patients with HBeAg-negative chronic liver disease from three different parts of China (Beijing, Shanghai and Hong Kong), 69% of genotype C patients in Hong Kong belonged to sub genotype Cs whereas 97% of genotype C HBV in Shanghai and Beijing belonged to subgenotype Ce ( $P < 0.0001$ ) [23].

31. Regarding HBV disease course, local studies suggested that patients infected with genotype C have a higher risk of cirrhosis and HCC development [21, 24], as well as more severe histological fibrosis [25]. A recent meta-analysis concluded that genotype C hepatitis B virus was associated with a higher risk of HCC than other major hepatitis B virus genotypes [26]. Among HBV genotype C, subgenotype Cs appears to carry a worse prognosis than subgenotype Ce [23]. In a local study by the Chinese University of Hong Kong, patients infected by subgenotype Cs had the lowest serum albumin and highest alanine aminotransferase levels compared with subgenotypes Ce and Ba. And, patients infected by subgenotype Cs also had more severe histological

necroinflammation than subgenotype Ce [23]. However, the meta-analysis did not find significant difference in the risk of HCC between HBV-infected patients with subgenotype Ce and Cs [26].

32. Nevertheless, in a study of end-stage HBV-related liver disease patients requiring transplantation, those with genotype B had significantly more pre-transplant acute flare and worse liver function while genotype C patients had a greater risk and severity of recurrence due to lamivudine-resistant mutants [27].

33. In a case control study, it was concluded that HCC patients had a significantly higher prevalence of core promoter mutations and genotype C but the association with HCC is mediated via the former [28]. A study of 5080 chronic HBV patients focusing on familial HCC found 22 such families, giving a prevalence of 4.3 families/1000 HBV carriers [29]. Age of onset of HCC is significantly younger in familial HCC than sporadic cases, and it progressively decreased down the generations, suggesting an anticipation phenomenon.

### **Hepatitis B Vaccination**

34. Occurrence of new HBV infection is dependent on the interplay of multiple factors, including size of HBV pool, proportion of susceptible population and chance of exposure to the virus. The universal vaccination programme for newborns, increased vaccination coverage in adults, practice of universal precaution in health care settings, screening of blood donors and promotion of safer sex all contributed to the reduced HBV incidence in Hong Kong. [30].

35. A 16-year follow up study of 1112 neonates born to HBsAg-positive mothers who received HBV vaccine and hepatitis B immunoglobulin at different schedules demonstrated the long term protective efficacy of immunisation [31]. Upon completion of the vaccination schedules, 92.6% developed antibody against surface antigen (anti-HBs) seroconversion. Thirty-nine (3.5%) babies were tested positive for HBsAg and had become chronic carriers, 35 of which (89.7%) occurred before one year of age. At the end of the 16<sup>th</sup> year, 610 subjects (54.9%) returned for blood test evaluation. Although the anti-HBs seroconversion rate dropped to 33.3% at the 16<sup>th</sup> year and a total of 90 (8%) vaccinees developed anti-HBc seroconversion, none was found to have breakthrough infection to become chronic HBV infection. At the 30<sup>th</sup> year of follow-up, 246 (22.1%) vaccinees returned for blood tests [32]. The anti-HBs

seroconversion rate was maintained at 37.4% at the 30<sup>th</sup> year. Although two and one subjects developed anti-HBc seroconversion at the 21<sup>st</sup> and 25<sup>th</sup> year respectively, there was no new development of HBsAg positivity detected. These findings demonstrated the long-term protective efficacy of neonatal hepatitis B immunization among high risk individuals up to at least 30 years. In another study comparing three different HBV vaccine regimens without boosters given to 318 HBV negative children recruited at age 3 months to 11 years and followed up annually, no subjects tested positive for HBsAg up to 18 years of follow up (88 subjects). A total of 88 anamnestic responses with significant increase in anti-HBs titers were documented in 70 subjects; 3 subjects had benign breakthrough HBV infection with isolated anti-HBc seroconversion [33].

36. Universal neonatal HBV vaccination programme has been in place in Hong Kong since 1988. The coverage rate for the birth dose of HBV vaccine among infants born locally from 2010 to 2017 was consistently above 99% (unpublished DH data). There is generally a slight decline in the coverage rate for the second or the third dose. The drop may be related to two factors: some local-births have returned to Mainland after delivery and did not attend MCHC for services, and some babies received the vaccine in the private sector instead of MCHC.

37. DH has been conducting immunisation coverage surveys (ICS) every two or three years starting from 2001 to determine immunisation the coverage rates of all vaccines, including HBV vaccination among children aged 2 to 5 years and attending pre-primary institutions including kindergartens and child care centers. Results from ICS conducted in 2001, 2003, 2006, 2009 and 2012 confirmed high coverage rates of hepatitis B vaccination [34, 35, 36]. In the latest round of ICS conducted in 2015 (unpublished DH data), a total of 8723 children enrolled in 51 pre-primary institutions participated in the survey, reaching an overall response rate of 81.3%. Similar to previous years, the 2015 survey demonstrated a satisfactorily high coverage rate of HBV vaccination (Box 44).

38. Apart from universal neonatal HBV vaccination programme, supplementary Primary 6 vaccination programme was introduced in 1998. The coverage rate for three doses of HBV vaccine had been consistently above 99% in the past decade but showed a slight decline in 2015/16 to 97.9% for the third dose. Of note, this coincided with a change of survey methodology in 2015 and an underestimation of the actual coverage was possible (Box 45). With a high coverage of the neonatal HBV



vaccination programme, the number of Primary 6 students eligible for HBV vaccination continued to decrease in the past decade (from 17 171 in 2000/01 to 707 in 2016/17). The number of students who did not receive the third dose vaccination remained stable at a few hundred per year.

39. In 2009, a HBsAg seroprevalence study was conducted among 1913 children aged 12 to 15 years (unpublished DH data). The study found an HBsAg seroprevalence of 0.78% (95% confidence interval 0.39-1.16%, [Box 46](#)) in these children who were born after the implementation of universal neonatal HBV vaccination programme. This result showed that Hong Kong had already achieved a time-bound goal of reducing chronic HBV infection rate to less than 2% among 5 year-old children by the year of 2012, as set by the Western Pacific Regional Office (WPRO) of the WHO. In July 2011, Hong Kong was verified by WPRO as having successfully achieved the goal of HBV control. Based on the same study findings, Hong Kong was also verified as of June 2013 as having met the goal of achieving a seroprevalence of less than 1%.

40. In the CRPVH 2001 study, about 16% of the telephone-interviewed subjects reported a history of HBV vaccination, with a higher frequency in persons below 50 years of age. Some 83% of them reported having completed the vaccination course. Over 99% had the cost paid by them or borne by their employers. In another recent local survey by face-to-face questionnaire interview on over 1900 adult Chinese, fifty-eight percent (n=1151) of the subjects had been tested for HBV during adulthood. Among those tested negative for HBV infection, fifty-eight percent (n=506) of them reported subsequent HBV vaccination [37]. Age, occupation, having children, and family monthly income, were independent factors associated with vaccination in the study. Overall, the persistent significant level of HBsAg seroprevalence in the local population, though declining, means a significant disease burden in the years to come.

### **Current Situation of Hepatitis C**

41. From 2002 to 2017, a total of 127 cases of acute hepatitis C infection were reported to DH under the statutory notification system ([Box 1](#)), with one to fourteen cases reported annually from 2002 to 2015, and a record high of 39 cases in 2016. A review by the Centre for Health Protection entitled “Hepatitis C in Hong Kong, 2008 to 2011” [38] showed that among the 22 laboratory confirmed acute hepatitis C cases reported to DH from January 2008 to October 2011, there were 17 males and 5

females, most (86%) acquired the infection locally. The median age was 47.5 years. Majority (86%) was ethnic Chinese. Five (23%) of them reported history of injecting drug use while no particular risk factor was identified for the remaining cases.

42. Of the 39 cases in 2016, 31 were male (79%), with age ranged from 23 to 94 years (median: 42 years). Thirteen (33%) required hospitalisation and no fatalities were recorded. With regard to the potential risk exposures, one case reported having tattoo procedure and two cases were identified as injecting drug users. Two cases reported to have sexual partners who were HCV carriers. Among the 31 male cases reported, 23 (74%) were known men who have sex with men. There was also one case who had history of repeated hospital admissions and had received multiple transfusions of blood product during the incubation period. Epidemiological investigation and contact tracing did not identify other acute hepatitis C cases and the source of infection in this case could not be determined. For the rest of the cases, no epidemiological linkage was identified and all cases were regarded as sporadic. There have been overseas reports of rising incidence of sexual transmission of HCV among men who have sex with men [39]. Further study and monitoring is required of the possibility that this is also the case for Hong Kong.

43. Although HCV shares similar transmission routes with hepatitis B, the epidemiology of two infections are different in Hong Kong. While HBV is still prevalent in many populations in Hong Kong, HCV prevails only in isolated communities.

44. Data from new blood donors who were mostly adolescents and young adults in the last decade suggested that HCV prevalence was around 0.1% locally, with the figure in 2017 being 0.06% (95% confidence interval; 0.03 - 0.08%) (Box 47). Findings of the household study of the entire spectrum of adult age groups conducted in 2001 further supported the uncommon occurrence of HCV infection among general population in Hong Kong; the overall positive rate was 0.3% in 936 subjects (95% confidence interval, 0.07%-0.94%) (Box 49). From 1999 to 2016, 10 of 2294 (0.4%) clients who attended the Therapeutic Prevention Clinic (TPC) at Integrated Treatment Centre (ITC) of CHP, DH for post-exposure management were tested positive for anti-HCV. Nine (90%) cases were non-HCW and all cases were already HCV infected at time of injury (Box 50).

45. From studies published in the early 1990s, it was shown that anti-HCV was more commonly found in injecting drug users (IDU, 66.8%), haemophilia (56%), haemodialysis (4.6%) and other patients requiring frequent blood/blood product

transfusions but not persons at risk through sexual contact [40]. In a more recent analysis of HCV positive blood donors, of those with identifiable risk factors, history of blood transfusion (43.7%) was the most common risk factor, followed by intravenous drug use (34.9%) and tattoo (28.6%). The source of infection was unknown in more than half of the respondents in the study [41].

46. A survey in 2011 of haemophiliacs under local public care found 100 of 222 patients (45%) infected with hepatitis C [42]. Another study conducted for 51 haemodialysis patients found that 8 (16%) were positive for anti-HCV by second generation enzyme immunoassay and 1 (2%) for HCV RNA alone, giving an overall infection rate of 18% [43]. This study also found a new infection rate of 4.9% per patient-year upon longitudinal follow up of 19 months.

47. Injecting drug use has been an important route of HCV acquisition. Results of testing non-random samples from drug users under treatment showed a HCV positive rate of 74% in 1988/1989 and 46% in 2000/2001 (Box 51). An HCV seroprevalence study in 2006 conducted in methadone clinics targeting IDU echoed the high prevalence rate of HCV in this community [44]. Of 567 IDU participants recruited in 2006, the prevalence of anti-HCV was 85% (95% confidence interval 82.5 – 88.3%). Another study in 2011 involving 622 IDU recruited at their gathering places found a similar figure of 81.7% (95% confidence interval 78.6 - 84.7%) infected with HCV [45]. In this study, the majority (84.7%) were male with a median age of 53 years. The median heroin injection duration was 25 years. Injection duration, current or recent injection, ever sharing injecting equipment and concomitant use of other drugs e.g. midazolam were independent factors associated with HCV infection in the two studies. In the recent New Life New Liver Project, which provided targeted HCV screening and education to ex-IDU in the community, 56% of 234 subjects screened were HCV positive. The number needed to screen to detect one patient with positive HCV was 1.8 (95% confidence interval 1.6-2.0) [46].

48. HIV/AIDS patients, with a proportion being IDU, is another group with consistent data showing a comparatively high HCV prevalence (Box 52, 53). From 2000 to 2017, HCV/HIV coinfection among new patients attending ITC ranged from 1.5% to 24.8%. The decreasing trend of anti-HCV seroprevalence was largely attributed to the decreasing proportion of new patients acquiring HIV via injecting drug use. The prevalence rate appears to be higher in male than female patients, likely related to the differential risk of parenteral and blood product exposure (Box 52). While HCV

infection was present in 1.7–6.3% of HIV/AIDS patients infected due to sexual contact, HCV was nearly universal in patients infected through drug injection (Box 53). It should be noted that, among male patients who acquired HIV via heterosexual contact and tested anti-HCV positive, 60.0% (30 out of 50 subjects) had a past history of injecting drug use (Box 53). Among those heterosexual male HIV infected patients without history of injecting drug use, the prevalence of anti-HCV was 3%.

49. There has been overseas data supporting sexual transmission of HCV among HIV-infected men who have sex with men [47]. The anti-HCV prevalence of subjects who contracted HIV via homosexual or bisexual contact in the ITC HIV/AIDS patient cohort has remained below 2% from screening since 2005. However, this figure has shown an increasing trend since 2012 with the number of individuals with HCV/HIV coinfection at the time of HIV diagnosis rising from 16 (1.3%) in 2013, to 50 (2.1%) in 2017 (Box 53).

50. From July to November 2013, ITC identified seven cases of recent HCV infection in Chinese HIV-infected MSM [48]. Five of the seven cases were also diagnosed to have recent syphilis infection during the period. None of them had history of injecting drug use. Phylogenetic analyses revealed that all cases belonged to the same genotype (genotype 3) although investigation showed no apparent linkage on their sexual exposure. An analysis on HIV-infected MSM attending ITC who had HCV seroconversion in the period 1999-2013 was subsequently performed [49]. Fourteen (1.1%) patients seroconverted, with an overall incidence rate of 0.22 per 100 patient-years. The incidence rate increased from 0.13 per 100 patient-years before 2002 to 0.19 per 100 patient years in 2002-2007 and 0.47 per 100 patient-years in 2008-2013. Genotype 3 was most commonly detected. Compared with the non-seroconverters, the seroconverters were of higher education level and had prior history of sexually transmitted infection. The overall higher HCV prevalence, and the increasing incidence of HCV among HIV-positive MSM, coupled with the hastened liver disease progression in HIV-infected patients [50], would no doubt result in a unique HCV/HIV coinfection that demands further attention.

51. Since 2003, laboratory surveillance for HCV in Hong Kong was enhanced to monitor the trend of anti-HCV among selected population groups, including blood donors from HKRCBTS, and selected in-patients from the Princess Margaret Hospital (PMH) and Prince of Wales Hospital (PWH, joined since 2005). Some 180,000-260,000 new and repeated blood donors of HKRCBTS were tested for anti-HCV each

year, among which the prevalence was consistently low at less than 0.1% since 2003. Whereas among the selected hospital patients tested in the past eleven years, the overall anti-HCV prevalence was 2.3% (Box 54). Anti-HCV was most commonly found in drug users, of which 51.8% were found positive, followed by patients with history of blood transfusion at 9.0%. Overall, the male-to-female ratio of HCV positive subjects was about 2.3 to 1, with a mean age of 51.4 years old (Box 55).

### Genotypes of Hepatitis C

52. Genotypic studies in Hong Kong has identified that 1b and 6a were the prevalent HCV genotypes locally, a scenario different from that in western countries where 1a predominated [51]. In an early study of 212 blood donors tested anti-HCV positive from 1991 to 1994, the commonest genotype found was 1b (58.8%), followed by 6a (27.0%) [52]. In another study of hospitalized patients with HCV testing for clinical indications 1b was the commonest type found in patients with chronic liver diseases and chronic renal failure [53]. According to a local study of patients on renal replacement therapy, the predominant genotype was 1b, followed by 1a and 6a [54]. Yet, the commonest genotype in intravenous drug users was genotype 6. A retrospective analysis of 106 intravenous drug users and 949 non-drug users with samples collected between December 1998 and May 2004 also confirmed the significant high prevalence of genotype 6a in drug users (58.5%) followed by 1b (33.0%), in contrast to 63.6% for 1b and 23.6% for 6a in non-drug users [55]. Besides intravenous drug use, age and sex were independent factors associated with HCV genotypes in this study. In a methadone clinic-based study published in 2011, out of 273 IDUs with different periods of initiating injection, 52% had genotype 6a and 38% had 1b. Both genotypes 1b and 6a were prevalent among older injectors, while subtype 3a was more common in young injectors and those initiating injection more recently during 1995-2006. Moreover, phylogenetic analysis revealed no specific clustering of any subtype or genotype, which did not suggest any outbreak of HCV among the study population. The extensive use of methadone widely available since 1980s may have protected Hong Kong from the emergence of HCV clusters among injection drug users [56].

53. For the HIV-infected MSM attending ITC who were diagnosed with acute HCV infection between 2009 to 2014, genotype 3a was the most prevalent (63.6%), followed by 1a (18.2%) and 6a (9.1%). The high prevalence of genotype 3a in MSM was in stark contrast to its rarity among HCV-infected IDU in Hong Kong. Phylogenetic analyses revealed a monophyletic HCV-3a cluster with members all diagnosed

between 2013 and 2014, and a homologous pair with HCV-6a genotype. However, there was no temporal or genetic clustering of the corresponding HIV sequences [57].

54. The natural history of 138 HCV genotype 1 patients (median age: 50 years) was compared with that of 78 HCV genotype 6 patients (median age: 46.5 years) in Queen Mary Hospital [58]. Both genotypes share a similar natural history based on liver biochemistry, HCV viral load, and probability of cirrhotic complications and mortality after a median follow-up period of over 5 years.

### **Liver Cancer – Major Morbidity and Mortality from Viral Hepatitis**

55. Chronic HBV and HCV infection are important risk factors for cirrhosis and liver cancer. Globally 788 000 people died of liver cancer in 2015, and HBV and HCV accounted for approximately 80% of liver cancer cases [59]. Local studies showed that 75-80% of hepatocellular cancers in Hong Kong were related to chronic HBV infection, and 3-6% cases were related to chronic HCV infection. HBV and HCV co-infection accounted for another 0.4-3% [60]. Among 76 liver transplants performed in Queen Mary Hospital due to cirrhosis from 1999 to 2000, 51 and 7 were related to hepatitis B and C respectively [61].

56. Apart from chronic HBV and HCV infection, other risk factors for liver cancer include excessive alcohol consumption and consumption of aflatoxin contaminated food. In Hong Kong, the age-standardised incidence rate and death rate of liver cancer is higher in male.

57. According to the data from the Hong Kong Cancer Registry [62], liver cancer, including neoplasm of liver and intrahepatic bile ducts, was the fourth commonest cancer in men and tenth commonest cancer in women in 2016. There were 1810 new registered cases of liver cancer, with 1391 cases of males and 419 cases of females (male to female ratio was about 3.3 to 1). There was a downward trend for the age-standardized incidence rate for both male and female in the past decade (Box 57). The figures were 23 for male and 5.7 for female per 100 000 standard population in 2016.

58. In 2016, liver cancer was the third leading cause of cancer deaths in Hong Kong. There were 1540 registered mortality from liver cancer. There was a downward trend for the age-standardized mortality rate for both sexes in the past decade (Box 59). The figures were 18 for male and 4.9 for female per 100 000 standard population in 2016 [62].

# SURVEILLANCE OF VIRAL HEPATITIS IN HONG KONG

## 2017 UPDATE

### Acute viral hepatitis under the disease notification system

#### Tabulated results


Box	Title	Page
Box 1.	Number of cases of viral hepatitis reported to the Department of Health between 1974 and 2017 (Data source: CHP, DH)	24
Box 2.	Reported cases of viral hepatitis from 1966 to 2017 (Data source: CHP, DH)	25
Box 3.	Breakdown of viral hepatitis by aetiology reported from 1996 to 2017 (Data source: CHP, DH)	26
<a href="#">Hepatitis A</a>		
Box 4.	Number of hepatitis A cases reported from 2003 to 2017 (Data source: CHP, DH)	27
Box 5.	Sex distribution of hepatitis A cases reported from 2003 to 2017 (Data source: CHP, DH)	28
Box 6.	Age distribution by proportion of reported cases of hepatitis A, 1989-2017 (Data source: CHP, DH)	29
Box 7.	Notification rates and death rates of hepatitis A, 1988 - 2017 (Data source: CHP, DH)	30
<a href="#">Hepatitis B</a>		
Box 8.	Number of hepatitis B cases reported from 1995 to 2017 (Data source: CHP, DH)	31
Box 9.	Sex distribution of hepatitis B cases reported from 1995 to 2017 (Data source: CHP, DH)	32
Box 10.	Age distribution of hepatitis B cases reported from 1995 to 2017 (Data source: CHP, DH)	33
<a href="#">Hepatitis C</a>		
Box 11.	Number of hepatitis C cases reported from 2002 to 2017 (Data source: CHP, DH)	34
Box 12.	Sex distribution of hepatitis C cases reported from 2005 to 2017 (Data source: CHP, DH)	35
Box 13.	Age distribution of hepatitis C cases reported from 2005 to 2017 (Data source: CHP, DH)	36
<a href="#">Hepatitis E</a>		
Box 14.	Number of reported hepatitis E cases reported from 1996 to 2017 (Data source: CHP, DH)	37
Box 15.	Mean and median plot of reported cases of hepatitis E by month from 1997 to 2017 (Data source: CHP, DH)	38
Box 16.	Sex distribution of hepatitis E cases reported from 1996 to 2017 (Data source: CHP, DH)	39
Box 17.	Age distribution by proportion of reported cases of hepatitis E from 1996 to 2017 (Data source: CHP, DH)	40
Box 18.	Notification rates and death rates of hepatitis E from 1996 to 2017 (Data source: CHP, DH)	41

**Box 1. Number of cases of viral hepatitis reported to the Department of Health between 1974 and 2017 (Data source: CHP, DH)**

Year	A	B	NAN B	C	E	Un- classifie d	Hepatitis (not elsewhere classified)	Total
1974		<i>notifiable since 1974</i>						639
1975								1761
1976								969
1977								1008
1978								1230
1979								964
1980								1554
1981								1738
1982								1814
1983								1783
1984								1780
1985								1601
1986								1425
1987								1554
1988	1187	250	465			496		2398
1989	618	136	154			324		1232
1990	1362	178	183			261		1984
1991	1297	150	200			154		1801
1992	3626	157	301			273		4357
1993	874	116	203			80		1273
1994	557	112	125			41		835
1995	491	102	55			18		666
1996	264	144	-	-	11	-	58	477
1997	595	100	-	-	4	-	37	736
1998	474	145	-	-	16	-	29	664
1999	426	152	-	-	8	-	31	617
2000	505	137	-	-	11	-	30	683
2001	494	134	-	-	26	-	23	677
2002	267	121	-	4	28	-	10	430
2003	107	98	-	-	19	-	8	232
2004	121	134	-	1	38	-	6	300
2005	64	105	-	1	34	-	0	204
2006	76	123	-	2	34	-	0	235
2007	69	74	-	1	65	-	0	209
2008	71	83	-	3	90	-	-	247
2009	64	80	-	3	73	-	-	220
2010	65	73	-	11	118	-	-	267
2011	46	70	-	5	119	-	-	240
2012	43	47	-	3	150	-	-	243
2013	44	40	-	10	90	-	-	184
2014	46	41	-	12	93	-	-	192
2015	138	29	-	14	84	-	-	265
2016	98	37	-	39	96	-	-	270
2017	117	33	-	18	64	-	-	232



**Box 2. Reported cases of viral hepatitis from 1966 to 2017 (Data source: CHP, DH)**




**Case definition:**

Hepatitis A – positive for anti-HAV IgM  
 Hepatitis C – positive for anti-HCV


Hepatitis B – positive for anti-HBc IgM  
 Hepatitis E - positive for anti-HEV

**Box 3. Breakdown of viral hepatitis by aetiology reported from 1996 to 2017 (Data source: CHP, DH)**




## HEPATITIS A

Box 4. Number of hepatitis A cases reported from 2003 to 2017 (Data source: CHP, DH)




## HEPATITIS A

Box 5. Sex distribution of hepatitis A cases reported from 2003 to 2017 (Data source: CHP, DH)



# HEPATITIS A


Box 6. Age distribution by proportion of reported cases of hepatitis A, 1989-2017 (Data source: CHP, DH)



Year


# HEPATITIS A

Box 7. Notification rates and death rates of hepatitis A, 1988 - 2017 (Data source: CHP, DH)




## HEPATITIS B

Box 8. Number of hepatitis B cases reported from 1995 to 2017 (Data source: CHP, DH)



## HEPATITIS B


Box 9. Sex distribution of hepatitis B cases reported from 1995 to 2017 (Data source: CHP, DH)






## HEPATITIS B

Box 10. Age distribution of hepatitis B cases reported from 1995 to 2017 (Data source: CHP, DH)




## HEPATITIS C

Box 11. Number of hepatitis C cases reported from 2002 to 2017 (Data source: CHP, DH)




## HEPATITIS C

Box 12. Sex distribution of hepatitis C cases reported from 2005 to 2017 (Data source: CHP, DH)




## HEPATITIS C

Box 13. Age distribution of hepatitis C cases reported from 2005 to 2017 (Data source: CHP, DH)




## HEPATITIS E

Box 14. Number of reported hepatitis E cases reported from 1996 to 2017 (Data source: CHP, DH)




## HEPATITIS E

Box 15. Mean and median plot of reported cases of hepatitis E by month from 1997 to 2017 (Data source: CHP, DH)




## HEPATITIS E

Box 16. Sex distribution of hepatitis E cases reported from 1996 to 2017 (Data source: CHP, DH)



# HEPATITIS E


Box 17. Age distribution by proportion of reported cases of hepatitis E from 1996 to 2017 (Data source: CHP, DH)





# HEPATITIS E

**Box 18. Notification rates and death rates of hepatitis E from 1996 to 2017 (Data source: CHP, DH)**



### Seroprevalence of hepatitis A and hepatitis E

#### Tabulated results

<b>Box</b>	<b>Title</b>	<b>Page</b>
Box 19.	Prevalence of anti-HAV in a collection of studies/testing between 1978 and 2009 (Data sources: Multiple sources)	43
Box 20.	Prevalence of anti-HAV in participants of Community Research Project for Viral Hepatitis (CRPVH) 2001 (Data source: DH)	44
Box 21.	Prevalence of anti-HAV in individuals with blood collected for serological diagnosis of conditions unrelated to hepatitis (Data source: PHLSB, CHP, DH)	44
Box 22.	Prevalence of anti-HAV at baseline screening of HIV/AIDS patients attending ITC from Jul 2007 to 2017 (Data source: ITC, CHP, DH)	45
Box 23.	Prevalence of anti-HAV per HIV risk at baseline screening of HIV/AIDS patients attending ITC from Jul 2007 to 2017 (Data source: ITC, CHP, DH)	46
Box 24.	Prevalence of anti-HEV in participants of Community Research Project for Viral Hepatitis (CRPVH) 2001 (Data source: DH)	47

## HEPATITIS A

**Box 19. Prevalence of anti-HAV in a collection of studies/testing between 1978 and 2009 (Data sources: Multiple sources)**

Age groups	1978	1987	1989	1993^	1995	1996		1998	2000	2001	2001	2002	2003	2004	2005	2006	2007	2008	2009
0 – 20	12.9% (0 - 10) 44.8% (11 - 20)	5.3% (0 - 10) 17.1% (11 - 20)	6.8% (0 - 10) 11.2% (11 - 20)	59.4% (M) 53.3% (F)	8.3%	- (0 - 10) 7.0% (11 - 20)	6.1%	5.4%	9.3%	4.58%	- (0 - 10) 12.5% (11 - 20)	5.3%	10.3%	14.7%	15.4%	20.0%	14.3%	16.7%	25.0%
21 – 30	75.0%	53.8%	58.8%	59.4% (M) 53.3% (F)	11.3%	-	11.8%	7.6%	17.5%	13.2%	26.8%	12.6%	13.2%	21.0%	28.2%	25.8%	19.4%	26.3%	30.3%
31 – 40	82.9%	85.1%	83.5%	59.4% (M) 53.3% (F)	49.0%	-	37.7%	40.8%	35.0%	41.3%	53.2%	46.7%	52.4%	43.8%	35.7%	50.0%	37.5%	47.4%	36.4%
>40	91.1%	94.7%	91.1% (41 - 50) 93.9% (>50)	94.5% (M) 91.0% (F)	70.5%	-	58.6%	66.7%	60.0%	71.1%	88.3% (41 - 50) 97.7% (>50)	58.1%	100.0%	50.0%	72.7%	80.0%	62.5%	71.4%	26.7%
Data source	A	B	C	D	E	F	E	E	E	E	G	E	E	E	E	E	E	E	E

^Figure is the average of age 0 – 40

Data sources:


- A. Study on left-over sera of 362 subjects, by Tsang et al of the University of Hong Kong [6]
- B. Study on stored sera of 702 healthy subjects, by Chin et al of the University of Hong Kong [5]
- C. Study on 1028 serum samples collected from individuals attending a health exhibition, by Lim et al of Department of Health. [63]
- D. Seroprevalence results reported in the press by Lai et al of the University of Hong Kong. [64]
- E. Pre-vaccination screening on students and staff of City University of Hong Kong: 553 (1995), 669 (1996), 608 (1998), 395 (2000), 592 (2001), 371 (2002), students and staff of Baptist University of Hong Kong 240 (2001), 259 (2002), 153 (2003), 55 (2004), 77 (2005), 53 (2006), 54 (2007), 70(2008),63(2009) and students and staff of Lingnan University 125 (2003), 84 (2004). [Data from CHC-Group Medical Practice]
- F. Seroprevalence study in school children by Lee et al of the Chinese University of Hong Kong. [65]
- G. Community Research Project on Viral Hepatitis 2001. [2]

## HEPATITIS A

**Box 20. Prevalence of anti-HAV in participants of Community Research Project for Viral Hepatitis (CRPVH) 2001 (Data source: DH)**

Age group	No. Tested	Anti-HAV +ve (%)
18-29	137	27 (19.7%)
30-39	223	116 (52.0%)
40-49	291	248 (85.2%)
50-59	170	161 (94.7%)
60 & over	115	113 (98.3%)
All	936	665 (71.0%)

**Box 21. Prevalence of anti-HAV in individuals with blood collected for serological diagnosis of conditions unrelated to hepatitis (Data source: PHLBS, CHP, DH)**



Year	Age group (years)													
	0-10		11-20		21-30		31-40		41-50		51-60		>60	
	No. tested	%	No. tested	%	No. tested	%	No. tested	%	No. tested	%	No. tested	%	No. tested	%
2000	420	8	190	19	200	31	190	59	100	95	-	-	-	-
2005	200	8	181	18	187	35	200	54	100	83	100	98	-	-
2010	96	16	100	22	100	37	95	54	100	64	100	91	100	100
2015	160	49	162	49	122	53	127	51	99	59	70	86	58	97

## HEPATITIS A

**Box 22. Prevalence of anti-HAV at baseline screening of HIV/AIDS patients attending ITC from Jul 2007 to 2017 (Data source: ITC, CHP, DH)**

Year (No. of patients)	Age	No. tested	Anti-HAV +ve (%)
2007 Jul-Dec (n=308)	<20	0	0 (0.0%)
	20-29	64	28 (43.8%)
	30-39	202	89 (44.1%)
	40-49	30	17 (56.7%)
	>=50	12	10 (83.3%)
2008 (n=506)	<20	2	1 (50.0%)
	20-29	101	39 (38.6%)
	30-39	282	142 (50.4%)
	40-49	77	49 (63.6%)
	>=50	44	42 (95.5%)
2009 (n=228)	<20	2	0 (0.0%)
	20-29	58	23 (39.7%)
	30-39	91	43 (47.3%)
	40-49	52	31 (59.6%)
	>=50	25	23 (92.0%)
2010 (n=222)	<20	3	0 (0.0%)
	20-29	41	18 (43.9%)
	30-39	82	49 (59.8%)
	40-49	54	34 (63.0%)
	>=50	42	35 (83.3%)
2011 (n=208)	<20	2	0 (0.0%)
	20-29	45	18 (40.0%)
	30-39	57	29 (50.9%)
	40-49	66	44 (66.7%)
	>=50	38	34 (89.5%)
2012 (n=361)	<20	6	0 (0.0%)
	20-29	64	18 (28.1%)
	30-39	105	44 (41.9%)
	40-49	111	70 (63.1%)
	>=50	75	56 (74.7%)
2013 (n=432)	<20	5	2 (40.0%)
	20-29	90	21 (23.3%)
	30-39	102	44 (43.1%)
	40-49	112	65 (58.0%)
	>=50	123	107 (87.0%)
2014 (n=375)	<20	8	1 (12.5%)
	20-29	135	42 (31.1%)
	30-39	96	42 (43.8%)
	40-49	68	32 (47.1%)
	>=50	68	58 (85.3%)

## HEPATITIS A

Year (No. of patients)	Age	No. tested	Anti-HAV +ve (%)
2015 (n=378)	<20	13	6 (46.2%)
	20-29	113	31 (27.4%)
	30-39	118	54 (45.8%)
	40-49	69	43 (62.3%)
	>=50	65	60 (92.3%)
2016 (n=344)	<20	4	0 (0.0%)
	20-29	106	24 (22.6%)
	30-39	121	46 (38.0%)
	40-49	57	30 (52.6%)
	>=50	56	45 (80.4%)
2017 (n=394)	<20	10	4 (40.0%)
	20-29	115	30 (26.1%)
	30-39	109	43 (39.4%)
	40-49	74	45 (60.8%)
	>=50	86	67 (77.9%)

**Box 23. Prevalence of anti-HAV per HIV risk at baseline screening of HIV/AIDS patients attending ITC from Jul 2007 to 2017 (Data source: ITC, CHP, DH)**

HIV risk	No. tested	Anti-HAV +ve (%)
Heterosexual male	720	514 (71.4%)
Heterosexual female	463	351 (75.8%)
Homo/Bi-sexual	2316	859 (37.1%)
Drug user	193	165 (85.5%)
Blood/blood product recipient	24	18 (75.0%)
Perinatal	9	0 (0.0%)
Undetermined	31	21 (67.7%)
Total	3756	1928 (51.3%)

## HEPATITIS E

### Box 24. Prevalence of anti-HEV in participants of Community Research Project for Viral Hepatitis (CRPVH) 2001 (Data source: DH)

Age group	No. Tested	Anti-HEV +ve (%)
18-29	137	11 (8.0%)
30-39	222	32 (14.4%)
40-49	290	70 (24.1%)
50-59	170	39 (22.9%)
60 & over	115	24 (20.9%)
All	934	176 (18.8%)

# SURVEILLANCE OF VIRAL HEPATITIS IN HONG KONG

## 2017 UPDATE

### Seroprevalence and vaccination coverage of hepatitis B

#### Tabulated results


Box	Title	Page
Box 25.	Prevalence of HBsAg in new blood donors from 1990 to 2017 (Data source: HKRCBTS)	50
Box 26.	HBsAg prevalence and its sex and age breakdown in new blood donors in 2017 (Data source: HKRCBTS)	51
Box 27.	HBsAg prevalence among university students/staff (Data source: City University Health Centre (till 2002), Baptist University Health Centre (2001 to 2009) & Lingnan University Health Service (2003 and 2004))	51
Box 28.	HBsAg prevalence from the FPAHK's Clinical Services (Data source: FPA)	52
Box 29.	HBsAg prevalence in antenatal women from 1990 to 2017 (Data source: FHS and PHLSB, CHP, DH)	53
Box 30.	HBsAg prevalence and age breakdown of antenatal mothers from 1990 to 2017 (Data source: FHS, DH)	54
Box 31.	Number of antenatal mothers with positive HBsAg by age, from 1990 to 2017 (Data source: FHS, DH)	55
Box 32.	Prevalence of hepatitis B markers in police officers, by sex from 1996 to 2006 and 2012 to 2017 (Data source: DH)	56
Box 33.	Prevalence of hepatitis B markers in police officers, by age from 1996 to 2006 and 2012 to 2017 (Data source: DH)	58
Box 34.	Prevalence of HBsAg from the Community Research Project on Viral Hepatitis (CRPVH) 2001 (Data source: DH)	59
Box 35.	Prevalence of HBsAg in newly recruited health care workers of DH from 2001 to 2017 (Data source: DH)	60
Box 36.	HBsAg prevalence among tuberculosis patients treated at chest clinics from 2005 to 2015 (March to May) (Data source: TB and Chest Service, CHP, DH)	61
Box 37.	HBsAg prevalence, stratified by age, among tuberculosis patients treated at chest clinics from 2005 to 2015 (March to May) (Data source: TB and Chest Service, CHP, DH)	62
Box 38.	Prevalence of hepatitis B markers in persons attending Therapeutic Prevention Clinic of Integrated Treatment Centre (ITC) for post-exposure management, from July 1999 to 2016 (Data source: ITC, CHP, DH)	63
Box 39.	Prevalence of hepatitis B markers in drug users from 1990 to 2010 (Data source: PHLSB, CHP, DH)	64



Box 40.	Prevalence of HBsAg at baseline screening of HIV/AIDS patients attending ITC from 2000 to 2017 (Data source: ITC, CHP, DH)	65
Box 41.	Prevalence of HBV infection per HIV risk at baseline screening of HIV/AIDS patients attending ITC from 2000 to 2017 (Data source: ITC, CHP, DH)	66
Box 42.	HBsAg prevalence (%) in new blood donors, pre-marital screening and antenatal women from 1990 to 2017 (Data source: multiple sources)	67
Box 43.	HBsAg prevalence (%) in other selected populations from 1990 to 2017 (Data source: multiple sources)	68
Box 44.	Hepatitis B immunisation coverage rates among children aged 2 to 5 by year of birth (Data source: ref 34, 35, 36 & unpublished DH data)	69
Box 45.	Cumulative statistics of the supplementary hepatitis B vaccination programme for Primary 6 students from the school years 2000 to 2017 (Data source: DH)	70
Box 46.	HBsAg seroprevalence by age among children aged 12 to 15 years in 2009 (Data source: unpublished data of DH)	71

## HEPATITIS B

Box 25. Prevalence of HBsAg in new blood donors from 1990 to 2017 (Data source: HKRCBTS)



## HEPATITIS B

**Box 26. HBsAg prevalence and its sex and age breakdown in new blood donors in 2017 (Data source: HKRCBTS)**


Age Group	Male		Female	
	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)
16-19	6667	22 (0.3%)	8366	21 (0.3%)
20-29	3770	31 (0.8%)	4220	21 (0.5%)
30-39	1907	50 (2.6%)	2856	39 (1.4%)
40-49	1107	32 (2.9%)	2101	39 (1.9%)
>49	611	17 (2.8%)	1190	16 (1.3%)
Total	14062	152 (1.1%)	18733	136 (0.7%)

**Box 27. HBsAg prevalence among university students/staff (Data source: City University Health Centre (till 2002), Baptist University Health Centre (2001 to 2009) & Lingnan University Health Service (2003 and 2004))**

Year	Aged below 21		Aged 21 – 30		Aged < 30	
	Total no. of cases	HBsAg +ve (%)	Total no. of cases	HBsAg +ve (%)	Total no. of cases	HBsAg +ve (%)
1994	305	7 (2.3%)	830	29 (3.5%)	1135	36 (3.2%)
1995	324	10 (3.1%)	768	33 (4.3%)	1092	43 (3.9%)
1996	348	4 (1.1%)	762	30 (3.9%)	1110	34 (3.1%)
1998	371	5 (1.3)	608	21 (3.5%)	979	26 (2.7%)
2000	230	7 (3.0%)	391	12 (3.1%)	621	19 (3.1%)
2001	508	13 (2.6%)	814	28 (3.4%)	1322	41 (3.1%)
2002	266	10 (3.8%)	483	13 (2.7%)	749	23 (3.1%)
2003	121	5 (4.1%)	214	8 (3.7%)	335	13 (3.9%)
2004	114	3 (2.6%)	217	4 (1.8%)	331	7 (2.1%)
2005	57	1 (1.8%)	115	0 (0.0%)	172	1 (0.6%)
2006	26	3 (11.5%)	104	1 (1.0%)	130	4 (3.1%)
2007	16	0 (0.0%)	82	1 (1.2%)	98	1 (1.0%)
2008	18	0 (0.0%)	82	1 (1.2%)	100	1 (1.0%)
2009	8	0 (0.0%)	56	0 (0.0%)	64	0 (0.0%)

# HEPATITIS B


Box 28. HBsAg prevalence from the FPAHK's Clinical Services (Data source: FPA)



Note: 1990-2010 only contain pre-marital check up  
 Start from 2011 contain both pre-marital and pre-pregnancy check up

# HEPATITIS B

Box 29. HBsAg prevalence in antenatal women from 1990 to 2017 (Data source: FHS and PHLSB, CHP, DH)



## HEPATITIS B


**Box 30. HBsAg prevalence and age breakdown of antenatal mothers from 1990 to 2017 (Data source: FHS, DH)**

Year	No. tested (% HBsAg +ve) according to age group of antenatal mothers				
	<20*	20-24	25-29	30-34	>34
1990	1044 (10.3%)	4671 (13.4%)	15228 (10.7%)	7639 (12.6%)	2780 (12.9%)
1991	987 (10.7%)	4620 (10.7%)	13151 (10.4%)	8168 (11.5%)	3063 (11.8%)
1992	928 (9.6%)	5065 (11.4%)	13093 (10.6%)	8788 (10.6%)	3470 (11.7%)
1993	984 (9.0%)	5589 (10.5%)	12345 (10.3%)	9395 (11.6%)	3798 (11.0%)
1994	951 (7.8%)	5723 (9.8%)	11590 (9.7%)	10158 (10.6%)	3998 (10.4%)
1995	922 (8.4%)	4979 (9.7%)	10619 (9.6%)	10112 (9.8%)	4283 (10.3%)
1996	842 (7.8%)	4765 (10.3%)	10137 (9.5%)	9759 (9.5%)	5908 (10.6%)
1997	902 (7.1%)	4207 (9.3%)	8895 (9.6%)	7982 (9.3%)	3897 (9.3%)
1998	911 (5.8%)	3887 (9.2%)	8507 (9.3%)	7418 (8.8%)	3851 (9.3%)
1999	794 (7.7%)	3777 (8.6%)	8068 (9.3%)	7196 (8.2%)	3975 (9.3%)
2000	618 (6.8%)	2974 (10.1%)	6466 (9.5%)	5818 (8.0%)	3192 (8.7%)
2001	659 (7.3%)	3516 (9.5%)	8330 (10.1%)	6936 (8.3%)	3915 (9.0%)
2002	484 (5.0%)	2829 (9.7%)	9120 (9.7%)	6351 (8.5%)	3414 (8.1%)
2003	548 (4.9%)	2880 (9.9%)	7614 (9.4%)	6789 (8.3%)	3602 (8.2%)
2004	510 (6.1%)	2854 (8.4%)	7161 (8.9%)	7732 (8.6%)	3856 (8.1%)
2005	445 (3.4%)	2753 (8.9%)	6063 (9.5%)	7869 (8.6%)	4114 (7.4%)
2006	516 (4.8%)	2590 (8.0%)	6271 (8.7%)	8637 (8.6%)	4514 (8.4%)
2007	520 (4.0%)	2929 (8.4%)	7301 (9.3%)	10232 (8.7%)	5551 (7.5%)
2008	533 (3.2%)	2968 (8.0%)	7652 (8.6%)	10354 (8.8%)	5838 (8.0%)
2009	434 (3.2%)	2830 (8.7%)	7444 (9.3%)	10156 (7.9%)	6071 (7.7%)
2010	446 (2.2%)	2903 (8.0%)	7817 (8.5%)	10211 (7.9%)	6385 (7.6%)
2011	447 (2.5%)	2898 (6.5%)	9010 (8.1%)	12273 (7.3%)	7552 (7.5%)
2012	463 (2.6%)	2467 (4.4%)	8161 (7.5%)	12664 (7.2%)	7437 (7.1%)
2013	423 (5.0%)	2237 (4.1%)	7526 (6.8%)	12466 (6.7%)	7168 (7.3%)
2014	366 (0.8%)	2252 (2.8%)	7901 (6.3%)	13488 (6.4%)	7692 (6.9%)
2015	409 (1.0%)	2439 (2.6%)	8589 (4.7%)	14434 (6.2%)	8656 (6.8%)
2016	328 (2.1%)	2123 (2.0%)	7580 (4.1%)	13018 (5.7%)	7923 (6.6%)
2017	274 (0.7%)	1897 (2.2%)	6624 (3.5%)	11476 (5.5%)	7611 (6.3%)

\* Figures before year 2001 refer to age group 15-19; figures after year 2001 refer to age group <20



## HEPATITIS B

Box 31 Number of antenatal mothers with positive HBsAg by age, from 1990 to 2017 (Date source: FHS, DH)



## HEPATITIS B

Box 32. Prevalence of hepatitis B markers in police officers, by sex from 1996 to 2006 and 2012 to 2017 (Data source: DH)



Note: Data was not available from 2007-Feb 2012

\* For a period between Mar-Dec 2012



## HEPATITIS B

Year	Male			Female			All		
	No. tested	HBsAg +ve (%)	Anti-HBs +ve (%)	No. tested	HBsAg +ve (%)	Anti-HBs +ve (%)	No. tested	HBsAg +ve (%)	Anti-HBs +ve (%)
1996	2080	138 (6.6%)	740 (35.6%)	413	15 (3.6%)	113 (27.4%)	2493	153 (6.1%)	853 (34.2%)
1997	4227	346 (8.2%)	1489 (35.2%)	472	26 (5.5%)	152 (32.2%)	4699	372 (7.9%)	1641 (34.9%)
1998	2316	177 (7.6%)	678 (29.3%)	284	16 (5.6%)	74 (26.1%)	2600	193 (7.4%)	752 (28.9%)
1999	1399	93 (6.6%)	424 (30.3%)	322	17 (5.3%)	91 (28.3%)	1721	110 (6.4%)	515 (29.9%)
2000	1300	83 (6.4%)	395 (30.4%)	244	3 (1.2%)	65 (26.6%)	1544	86 (5.6%)	460 (29.8%)
2001	1058	69 (6.5%)	330 (31.2%)	221	6 (2.7%)	78 (35.3%)	1279	75 (5.9%)	408 (31.9%)
2002	1374	77 (5.6%)	416 (30.3%)	270	10 (3.7%)	81 (30%)	1644	87 (5.3%)	497 (30.2%)
2003	1415	69 (4.9%)	388 (27.4%)	259	8 (3.1%)	71 (27.4%)	1674	77 (4.6%)	459 (27.4%)
2004	1105	58 (5.2%)	361 (32.7%)	188	5 (2.7%)	79 (42%)	1293	63 (4.9%)	440 (34%)
2005	1613	68 (4.2%)	562 (34.8%)	323	13 (4.0%)	137 (42.4%)	1936	81 (4.2%)	699 (36.1%)
2006	195	9 (4.6%)	74 (37.9%)	44	2 (4.5%)	20 (45.5%)	239	11 (4.6%)	94 (39.3%)
2012*	1494	49 (3.3%)	635 (42.5%)	338	6 (1.8%)	165 (48.8%)	1832	55 (3.0%)	800 (43.7%)
2013	1812	52 (2.9%)	751 (41.4%)	506	13 (2.6%)	207 (40.9%)	2318	65 (2.8%)	958 (41.3%)
2014	2267	59 (2.6%)	847 (37.4%)	560	15 (2.7%)	230 (41.1%)	2827	74 (2.6%)	1077 (38.1%)
2015	2563	71 (2.8%)	972 (37.9%)	621	17 (2.7%)	263 (42.4%)	3184	88 (2.8%)	1235 (38.8%)
2016	2450	49 (2.0%)	796 (32.5%)	561	9 (1.6%)	191 (34.0%)	3011	58 (1.9%)	987 (32.8%)
2017	2477	36 (1.5%)	768 (31.0%)	706	9 (1.3%)	228 (32.3%)	3183	45 (1.4%)	996 (31.3%)

Note: Data was not available from 2007-Feb 2012

\* For a period between Mar-Dec 2012

## HEPATITIS B

Box 33. Prevalence of hepatitis B markers in police officers, by age from 1996 to 2006 and 2012 to 2017 (Data source: DH)

Year	Age group														
	≤20			21-30			31-40			41-50			>50		
	No. tested	% HBsAg +ve	% Anti-HBs +ve	No. tested	% HBsAg +ve	% Anti-HBs +ve	No. tested	% HBsAg +ve	% Anti-HBs +ve	No. tested	% HBsAg +ve	% Anti-HBs +ve	No. tested	% HBsAg +ve	% Anti-HBs +ve
1996	17	0.0	35.3	733	4.8	24.4	1155	6.8	32.9	544	5.9	49.6	44	18.2	40.9
1997	15	6.7	46.7	1494	6.1	25.4	2081	7.3	35.0	999	11.4	46.6	110	13.6	55.5
1998	387	5.9	20.7	969	5.5	25.0	828	8.3	30.8	356	12.4	40.4	60	6.7	51.7
1999	270	4.4	24.1	799	6.1	27.5	428	6.8	31.8	202	8.9	42.1	22	9.1	40.9
2000	72	4.2	22.2	746	6.4	24.3	460	4.3	31.3	242	5.8	44.6	24	4.2	45.8
2001	68	4.4	30.9	602	5.8	28.4	339	5.6	30.7	225	6.2	40.0	45	8.9	48.9
2002	145	4.8	29.7	697	4.9	25.3	443	3.6	29.6	307	9.1	37.5	52	3.8	61.5
2003	72	1.4	16.7	702	4.8	22.9	505	4.6	26.5	357	5.0	38.1	38	2.6	42.1
2004	8	0.0	37.5	466	5.2	35.6	441	3.4	28.6	321	5.9	39.6	57	8.8	31.6
2005	80	1.3	52.5	791	3.8	32.7	533	4.3	31.0	427	4.2	43.3	105	8.6	45.7
2006	0	-	-	39	0.0	51.3	86	5.8	36.0	90	4.4	36.7	24	8.3	41.7
2012*	267	0.7	20.2	1169	2.1	47.3	122	6.6	53.3	203	5.9	47.8	71	11.3	43.7
2013	393	0.0	24.4	1635	2.7	43.8	95	4.2	57.9	133	11.3	46.6	62	3.2	46.8
2014	456	0.7	24.8	1789	1.9	37.8	188	6.4	48.9	280	6.4	51.1	114	6.1	46.5
2015	455	0.9	24.8	2077	2.4	38.9	221	5.4	50.7	309	5.5	46.9	122	4.1	47.5
2016	428	0.5	17.3	2250	1.6	33.2	154	5.2	53.2	125	7.2	49.6	54	3.7	42.6
2017	391	0.5	21.2	2594	1.3	31.7	182	2.2	46.7	13	38.5	30.8	3	0.0	66.7

Note: Data was not available from 2007-Feb 2012

\* For a period between Mar-Dec 2012


## HEPATITIS B

**Box 34. Prevalence of HBsAg from the Community Research Project on Viral Hepatitis (CRPVH) 2001 (Data source: DH)**

Age Group	Male		Female		Total	
	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)
18-30	72	6 (8.3%)	87	6 (6.9%)	159	12 (7.5%)
31-40	93	5 (5.4%)	144	20 (13.9%)	237	25 (10.5%)
41-50	100	20 (20.0%)	183	10 (5.5%)	283	30 (10.6%)
51 & Over	111	8 (7.2%)	146	7 (4.8%)	257	15 (5.8%)
Total	376	39 (10.4%)	560	43 (7.7%)	936	82 (8.8%)

## HEPATITIS B


**Box 35. Prevalence of HBsAg in newly recruited health care workers of DH from 2001 to 2017 (Data source: DH)**



Year	Male		Female	
	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)
2001	440	27 (6.1%)	613	36 (5.9%)
2002	499	23 (4.6%)	730	38 (5.2%)
2003	373	20 (5.4%)	531	27 (5.1%)
2004	307	13 (4.2%)	644	37 (5.7%)
2005	396	22 (5.6%)	956	51 (5.3%)
2006	220	8 (3.6%)	449	25 (5.6%)
2007	204	8 (3.9%)	102	4 (3.9%)
2008	232	7 (3.0%)	187	9 (4.8%)
2009	226	14 (6.2%)	328	14 (4.3%)
2010	307	15 (4.9%)	239	10 (4.2%)
2011	370	12 (3.2%)	233	3 (1.3%)
2012	318	18 (5.7%)	377	12 (3.2%)
2013	282	8 (2.8%)	418	19 (4.5%)
2014	261	3 (1.1%)	370	13 (3.5%)
2015	324	8 (2.5%)	391	15 (3.8%)
2016	278	8 (2.9%)	409	16 (3.9%)
2017	291	7 (2.4%)	452	16 (3.5%)

## HEPATITIS B

**Box 36. HBsAg prevalence among tuberculosis patients treated at chest clinics from 2005 to 2015 (March to May) (Data source: TB and Chest Service, CHP, DH)**



Year	Male		Female		Total	
	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)
2005	442	52 (11.8%)	242	17 (7.0%)	684	69 (10.1%)
2006	821	97 (11.8%)	446	27 (6.1%)	1267	124 (9.8%)
2007	768	96 (12.5%)	420	29 (6.9%)	1188	125 (10.5%)
2008	648	62 (9.6%)	382	30 (7.9%)	1030	92 (8.9%)
2009	759	73 (9.6%)	438	30 (6.8%)	1197	103 (8.6%)
2010	669	64 (9.6%)	353	22 (6.2%)	1022	86 (8.4%)
2011	674	77 (11.4%)	382	29 (7.6%)	1056	106 (10.0%)
2012	651	59 (9.1%)	367	27 (7.4%)	1018	86 (8.4%)
2013	664	70 (10.5%)	369	25 (6.8%)	1033	95 (9.2%)
2014	598	60 (10.0%)	393	24 (6.1%)	991	84 (8.5%)
2015	560	56 (10.0%)	314	18 (5.7%)	874	74 (8.5%)

## HEPATITIS B

**Box 37. HBsAg prevalence, stratified by age, among tuberculosis patients treated at chest clinics from 2005 to 2015 (March to May) (Data source: TB and Chest Service, CHP, DH)**

Year	Age group									
	0-19		20-39		40-59		≥60		Total	
	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)
2005	31	1 (3.2%)	168	11 (6.5%)	204	34 (16.7%)	281	23 (8.2%)	684	69 (10.1%)
2006	47	2 (4.3%)	314	21 (6.7%)	402	57 (14.2%)	504	44 (8.7%)	1267	124 (9.8%)
2007	57	1 (1.8%)	287	20 (7.0%)	374	60 (16.0%)	470	44 (9.4%)	1188	125 (10.5%)
2008	26	1 (3.8%)	256	14 (5.5%)	316	42 (13.3%)	432	35 (8.1%)	1030	92 (8.9%)
2009	45	0 (0.0%)	275	22 (8.0%)	370	56 (15.1%)	507	25 (4.9%)	1197	103 (8.6%)
2010	34	0 (0.0%)	224	15 (6.7%)	315	39 (12.4%)	449	32 (7.1%)	1022	86 (8.4%)
2011	35	0 (0.0%)	259	18 (6.9%)	303	45 (14.9%)	459	43 (9.4%)	1056	106 (10.0%)
2012	32	0 (0.0%)	261	21 (8.0%)	315	32 (10.2%)	410	33 (8.0%)	1018	86 (8.4%)
2013	54	1 (1.9%)	228	13 (5.7%)	320	41 (12.8%)	431	40 (9.3%)	1033	95 (9.2%)
2014	34	1 (2.9%)	211	8 (3.8%)	313	36 (11.5%)	433	39 (9.0%)	991	84 (8.5%)
2015	30	0 (0.0%)	187	13 (7.0%)	260	26 (10.0%)	397	35 (8.8%)	874	74 (8.5%)

## HEPATITIS B

**Box 38. Prevalence of hepatitis B markers in persons attending Therapeutic Prevention Clinic of Integrated Treatment Centre (ITC) for post-exposure management, from July 1999 to 2016 (Data source: ITC, CHP, DH)**

Year	Health care workers			Non- Health care workers			Total		
	No. tested	HBsAg +ve (%)	Anti-HBs +ve (%)	No. tested	HBsAg +ve (%)	Anti-HBs +ve (%)	No. tested	HBsAg +ve (%)	Anti-HBs +ve (%)
Jul-Dec 1999	23	2 (8.7%)	11 (47.8%)	87	13 (14.9%)	41 (47.1%)	110	15 (13.6%)	52 (47.3%)
2000	77	5 (6.5%)	56 (72.7%)	217	20 (9.2%)	91 (41.9%)	294	25 (8.5%)	147 (50.0%)
2001	103	2 (1.9%)	78 (75.7%)	313	20 (6.4%)	143 (45.7%)	416	22 (5.3%)	221 (53.1%)
2002	99	9 (9.1%)	62 (62.6%)	252	22 (8.7%)	133 (52.8%)	351	31 (8.8%)	195 (55.6%)
2003	96	6 (6.3%)	66 (68.8%)	201	24 (11.9%)	81 (40.3%)	297	30 (10.1%)	147 (49.5%)
2004	66	4 (6.1%)	41 (62.1%)	182	15 (8.2%)	97 (53.3%)	248	19 (7.7%)	138 (55.6%)
2005	49	3 (6.1%)	31 (63.3%)	206	13 (6.3%)	99 (48.1%)	255	16 (6.3%)	130 (51.0%)
2006	54	6 (11.1%)	33 (61.1%)	289	15 (5.2%)	151 (52.2%)	343	21 (6.1%)	184 (53.6%)
2007	54	1 (1.9%)	45 (83.3%)	228	18 (7.9%)	88 (38.6%)	282	19 (6.7%)	133 (47.2%)
2008	54	2 (3.7%)	39 (72.2%)	235	20 (8.5%)	111 (47.2%)	289	22 (7.6%)	150 (51.9%)
2009	56	1 (1.8%)	41 (73.2%)	297	22 (7.4%)	138 (46.5%)	353	23 (6.5%)	179 (50.7%)
2010	47	1 (2.1%)	33 (70.2%)	245	10 (4.1%)	137 (55.9%)	292	11 (3.8%)	170 (58.2%)
2011	54	1 (1.9%)	35 (64.8%)	270	12 (4.4%)	159 (58.9%)	324	13 (4.0%)	194(59.9%)
2012	70	2 (2.9%)	54 (77.1%)	311	16 (5.1%)	173 (55.6%)	381	18 (4.7%)	227 (59.6%)
2013	82	1 (1.2%)	64 (78.0%)	313	15 (4.8%)	149 (47.6%)	395	16 (4.1%)	213 (53.9%)
2014	79	3 (3.8%)	58 (73.4%)	330	9 (2.7%)	180 (54.5%)	409	12 (2.9%)	238 (58.2%)
2015	85	1 (1.2%)	66 (77.6%)	311	10 (3.2%)	172 (55.3%)	396	11 (2.8%)	238 (60.1%)
2016	118	2 (1.7%)	82 (69.5%)	343	12 (3.5%)	155 (45.2%)	461	14 (3.0%)	237 (51.4%)
Total	1266	52 (4.1%)	895 (70.7%)	4630	286 (6.2%)	2298 (49.6%)	5896	338 (5.7%)	3193 (54.2%)

## HEPATITIS B

**Box 39. Prevalence of hepatitis B markers in drug users from 1990 to 2010 (Data source: PHL SB, CHP, DH)**


Year	No. tested	HBsAg (%+ve)	Anti-HBs (%+ve)	Anti-HBc* (%+ve)	Any marker (%+ve)
1990	1067	13.4	59.0	15.7	90.8
1991	1517	14.4	54.4	20.5	89.3
1992	832	13.9	49.0	21.4	84.4
1993	744	14.4	43.4	16.4	69.2
1994	607	12.9	38.1	13.5	64.1
1995	190	10.5	36.8	12.1	58.9
1996	358	8.7	43.0	12.6	62.8
1997	290	6.6	36.2	15.9	53.4
1998	290	10.0	43.4	7.9	59.3
1999	725	11.2	44.8	13.8	67.2
2000	892	11.4	42.5	15.8	67.8
2001	654	11.6	41.3	17.3	70.2
2002	553	12.7	43.0	16.6	72.3
2003	198	10.1	42.4	12.6	65.2
2004	45	11.1	57.8	4.4	73.3
2005	26	11.5	46.2	11.5	69.2
2006	6	33.3	50.0	16.7	100.0
2007	11	0.0	81.8	9.1	90.9
2008	7	28.6	28.6	14.3	71.4
2009	11	9.1	72.7	9.1	100.0
2010	12	8.3	58.3	8.3	100.0

\*Anti-HBc was not tested in specimens that were HBsAg positive



## HEPATITIS B

**Box 40. Prevalence of HBsAg at baseline screening of HIV/AIDS patients attending ITC from 2000 to 2017 (Data source: ITC, CHP, DH)**




Year	Male		Female		Total	
	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)	No. tested	HBsAg +ve (%)
2000	57	6 (10.5%)	17	1 (5.9%)	74	7 (9.5%)
2001	75	11 (14.7%)	23	1 (4.3%)	98	12 (12.2%)
2002	112	14 (12.5%)	22	1 (4.5%)	134	15 (11.2%)
2003	93	12 (12.9%)	15	2 (13.3%)	108	14 (13.0%)
2004	115	20 (17.4%)	23	2 (8.7%)	138	22 (15.9%)
2005	132	8 (6.1%)	29	1 (3.4%)	161	9 (5.6%)
2006	188	26 (13.8%)	22	3 (13.6%)	210	29 (13.8%)
2007	216	27 (12.5%)	27	1 (3.7%)	243	28 (11.5%)
2008	203	22 (10.8%)	33	1 (3.0%)	236	23 (9.7%)
2009	170	16 (9.4%)	27	1 (3.7%)	197	17 (8.6%)
2010	160	20 (12.5%)	34	2 (5.9%)	194	22 (11.3%)
2011	167	17 (10.2%)	33	2 (6.1%)	200	19 (9.5%)
2012	226	27 (11.9%)	44	2 (4.5%)	270	29 (10.7%)
2013	263	15 (5.7%)	41	2 (4.9%)	304	17 (5.6%)
2014	301	24 (8.0%)	31	1 (3.2%)	332	25 (7.5%)
2015	328	20 (6.1%)	26	0 (0.0%)	354	20 (5.6%)
2016	304	22 (7.2%)	25	3 (12.0%)	329	25 (7.6%)
2017	326	28 (8.6%)	33	1 (3.0%)	359	29 (8.1%)

## HEPATITIS B

**Box 41. Prevalence of HBV infection per HIV risk at baseline screening of HIV/AIDS patients attending ITC from 2000 to 2017 (Data source: ITC, CHP, DH)**


HIV risk	No. tested	HBsAg +ve (%)	Anti-HBs +ve (%)
Heterosexual male	797	93 (11.7%)	371 (46.5%)
Heterosexual female	472	27 (5.7%)	202 (42.8%)
Homo/Bi-sexual	2356	198 (8.4%)	1267 (53.8%)
Drug user	259	41 (15.8%)	123 (47.5%)
Blood/blood product recipient	14	0 (0%)	5 (35.7%)
Perinatal	9	0 (0%)	2 (22.2%)
Undetermined	34	3 (8.8%)	14 (41.2%)
<b>Total</b>	<b>3941</b>	<b>362 (9.2%)</b>	<b>1984 (50.3%)</b>



## HEPATITIS B

Box 42. HBsAg prevalence (%) in new blood donors, pre-marital screening and antenatal women from 1990 to 2017 (Data source: multiple sources)

Year	New blood donors	Pre-marital screening	Antenatal women
1990	8.0	9.6	11.3
1991	8.0	9.6	10.9
1992	7.4	9.3	10.8
1993	6.7	8.7	10.1
1994	5.9	7.3	10.0
1995	6.0	7.9	9.7
1996	5.6	7.9	9.7
1997	5.2	7.6	9.3
1998	4.9	6.8	9.0
1999	4.4	6.7	8.8
2000	4.2	5.6	8.9
2001	4.0	5.1	9.2
2002	3.6	6.9	9.0
2003	3.2	7.1	8.8
2004	2.9	7.4	8.5
2005	2.6	6.8	8.6
2006	2.2	7.4	8.4
2007	1.8	7.1	8.5
2008	1.8	6.4	8.4
2009	1.6	6.9	8.2
2010	1.2	6.5	7.9
2011	1.1	6.4	7.4
2012	1.1	6.9	7.0
2013	1.1	6.3	6.6
2014	0.8	5.5	6.2
2015	1.0	5.3	5.7
2016	0.8	6.2	5.2
2017	0.9	4.8	5.0



## HEPATITIS B

Box 43 HBsAg prevalence (%) in other selected populations from 1990 to 2017 (Data source: multiple sources)

Year	University students/staff (aged 21-30)	Police officers	Health care workers	Drug users	Female sex workers	HIV/AIDS patients	Tuberculosis patients	TPC patients
1990	-	-	-	13.4	-	-	-	-
1991	-	-	6.2	14.4	-	-	-	-
1992	-	-	-	13.9	-	-	-	-
1993	-	-	4.4	14.4	-	-	-	-
1994	3.5	-	-	12.9	-	-	-	-
1995	4.3	-	7.0	10.5	6.8^	-	-	-
1996	3.9	6.1	4.2	8.7	6.8^	-	-	-
1997	-	7.9	-	6.6	6.8^	-	-	-
1998	3.5	7.4	-	10.0	6.8^	-	-	-
1999	-	6.4	2.2	11.2	-	-	-	13.6*
2000	3.1	5.6	5.4	11.4	-	9.5	-	8.5
2001	3.4	5.9	6.0	11.6	-	12.2	-	5.3
2002	2.7	5.3	5.0	12.7	-	11.2	-	8.8
2003	3.7	4.6	5.2	10.1	-	13	-	10.1
2004	1.8	4.9	5.3	-	-	15.9	-	7.7
2005	-	4.2	5.4	-	-	5.6	10.1	6.3
2006	1.0	4.6	4.9	-	-	13.8	9.8	6.1
2007	1.2	-	3.9	-	10.4**	11.5	10.5	6.7
2008	1.2	-	3.8	-	9.0	9.7	8.9	7.6
2009	0.0	-	5.1	-	6.5	8.6	8.6	6.5
2010	-	-	4.6	-	5.0	11.3	8.4	3.8
2011	-	-	2.5	-	7.2***	9.5	10.0	4.0
2012	-	3.0****	4.3	-	-	10.7	8.4	4.7
2013	-	2.8	3.9	-	-	5.6	9.2	4.1
2014	-	2.6	2.5	-	-	7.5	8.5	2.9
2015	-	2.8	3.2	-	-	5.6	8.5	2.8
2016	-	1.9	3.5	-	-	7.6	-	3.0
2017	-	1.4	3.1	-	-	8.1	-	-

\*For a period between Jul-Dec 1999; \*\*For a period between Aug-Dec 2007, \*\*\* For a period between Jan-Jul 2011, \*\*\*\* For a period between Mar-Dec 2012

^Figure is the average of 1995-1998

## HEPATITIS B

**Box 44. Hepatitis B immunisation coverage rates among children aged 2 to 5 by year of birth (Data source: ref 34, 35, 36 & unpublished DH data)**

Year of Survey	Year of Birth	First dose (%)	Second dose (%)	Third dose (%)
2001	1995	99.5	99.5	99.1
	1996	99.1	99	98.6
2003	1997	99.5	99.3	99.1
	1998	99.9	99.9	99.6
	1999	100	100	99.7
2006	2000	99.9	99.8	99.6
	2001	99.9	99.9	99.6
	2002	99.9	99.8	99.5
2009	2003	99.9	99.8	99.5
	2004	99.9	99.9	99.8
	2005	99.7	99.7	99.5
	2006	100	100	99.7
2012	2006	99.6	99.5	99.0
	2007	99.8	99.8	99.3
	2008	99.8	99.8	99.3
	2009	100	100	98.8
2015	2009	99.7	99.6	99.2
	2010	99.6	99.6	99.2
	2011	99.6	99.5	99.2
	2012	100	100	99.2


## HEPATITIS B

**Box 45. Cumulative statistics of the supplementary hepatitis B vaccination programme for Primary 6 students from the school years 2000 to 2017 (Data source: DH)**

	2000-2001	2001-2002	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014	2014-2015	2015-2016	2016-2017
Cumulative no. of Primary 6 students	85612	86052	86515	86208	83974	83164	81818	77273	73757	67310	63332	63394	57487	54845	52013	51009	52843
<b>First Dose</b>																	
Cumulative no. eligible for vaccination	17171	15479	14245	10625	8433	6648	6351	6204	5165	4698	3736	2509	2376	1992	1797	982	707
Cumulative no. administered	16985	15333	14084	10519	8313	6591	6262	6095	5043	4520	3563	2318	2237	1810	1606	729	542
Acceptance rate (at the present campaign)	98.90%	99.10%	98.90%	99.00%	98.60%	99.10%	98.60%	98.20%	97.60%	96.2%	95.4%	92.4%	94.1%	90.9%	89.4%	74.2%	76.7%
Coverage rate (for the whole Primary 6 population)	99.80%	99.80%	99.80%	99.90%	99.80%	99.90%	99.90%	99.90%	99.80%	99.7%	99.7%	99.7%	99.8%	99.7%	99.6%	98.4%	98.6%
<b>Second Dose</b>																	
Cumulative no. eligible for vaccination	17182	15485	14250	10626	8545	6710	6392	6243	5165	4698	3787	2573	2432	2033	1825	1025	749
Cumulative no. administered	16890	15206	13800	10341	8185	6573	6278	6068	4969	4398	3516	2286	2203	1718	1578	675	534
Acceptance rate (at the present campaign)	98.30%	98.20%	96.80%	97.30%	95.80%	98.00%	98.20%	97.20%	96.20%	93.6%	92.8%	88.8%	90.6%	84.5%	86.5%	65.9%	71.3%
Coverage rate (for the whole Primary 6 population)	99.70%	99.70%	99.50%	99.70%	99.60%	99.80%	99.80%	99.80%	99.70%	99.5%	99.6%	99.5%	99.6%	99.4%	99.5%	98.2%	98.5%
<b>Third Dose</b>																	
Cumulative no. eligible for vaccination	17771	16119	14918	11222	9300	7397	6986	6741	5575	5032	4104	2825	2692	2283	2096	1307	1065
Cumulative no. administered	16741	14947	13999	10069	8478	6965	6607	6273	4817	4409	*3526	2344	2232	1777	1708	835	771
Acceptance rate (at the present campaign)	94.20%	92.70%	93.80%	89.70%	91.20%	94.20%	94.60%	93.10%	86.40%	87.6%	85.9%	83.0%	82.9%	77.8%	81.5%	63.9%	72.4%
Coverage rate (for the whole Primary 6 population)	98.80%	98.60%	98.90%	98.70%	99.00%	99.50%	99.50%	99.40%	99.00%	99.1%	99.1%	99.2%	99.2%	99.1%	99.3%	97.9%	98.3%

## HEPATITIS B

Box 46. HBsAg seroprevalence by age among children aged 12 to 15 years in 2009 (Data source: unpublished data of DH)



### Seroprevalence of hepatitis C


#### Tabulated results

Box	Title	Page
Box 47.	Anti-HCV prevalence in new blood donors, 1991 to 2017 (Data source: HKRCBTS)	73
Box 48.	Anti-HCV prevalence and its sex and age breakdown in new blood donors in 2017 (Data source: HKRCBTS)	74
Box 49.	Prevalence of anti-HCV in participants of Community Research Project on Viral Hepatitis (CRPVH) 2001 (Data source: DH)	75
Box 50.	Prevalence of anti-HCV at baseline screening of injured persons attending Therapeutic Prevention Clinic of Integrated Treatment Centre (ITC), from July 1999 to 2016 (Data source: ITC, CHP, DH)	75
Box 51.	Prevalence of anti-HCV in drug users on rehabilitation (Data source: PHLSB, CHP, DH)	76
Box 52.	Prevalence of anti-HCV at baseline screening of HIV/AIDS patients attending ITC from 2000 to 2017 (Data source: ITC, CHP, DH)	76
Box 53.	Prevalence of anti-HCV per HIV risk at baseline screening of HIV/AIDS patients attending ITC from 2000 to 2017 (Data source: ITC, CHP, DH)	77
Box 54.	Prevalence of anti-HCV from screening of blood donors and clinical testing of patients in 2 hospital clusters under Hospital Authority from 2007 to 2017 (Data source: HKRCBTS, PMH Microbiology Laboratory, PWH Microbiology Laboratory)	78
Box 55.	Characteristics of anti-HCV positive subjects detected at HKRCBTS and 2 hospital clusters under Hospital Authority from 2005 to 2017 (Data source: HKRCBTS, PMH Microbiology Laboratory, PWH Microbiology Laboratory)	79



# HEPATITIS C

Box 47. Anti-HCV prevalence in new blood donors, 1991 to 2017 (Data source: HKRCBTS)



## HEPATITIS C

**Box 48. Anti-HCV prevalence and its sex and age breakdown in new blood donors in 2017 (Data source: HKRCBTS)**

Age Group	Male		Female	
	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)
16-19	6667	1 (0.01%)	8366	1 (0.01%)
20-29	3770	1 (0.03%)	4220	0 (0.00%)
30-39	1907	6 (0.31%)	2856	2 (0.07%)
40-49	1107	2 (0.18%)	2101	3 (0.14%)
>49	611	2 (0.33%)	1190	1 (0.08%)
Total	14062	12 (0.09%)	18733	7 (0.04%)

## HEPATITIS C

**Box 49. Prevalence of anti-HCV in participants of Community Research Project on Viral Hepatitis (CRPVH) 2001 (Data source: DH)**

Age group	No. Tested	Anti-HCV +ve (%)
18-29	137	0 (0.0%)
30-39	223	1 (0.4%)
40-49	291	0 (0.0%)
50-59	170	2 (1.2%)
60 & over	115	0 (0.0%)
All	936	3 (0.3%)

**Box 50. Prevalence of anti-HCV at baseline screening of injured persons attending Therapeutic Prevention Clinic of Integrated Treatment Centre (ITC), from July 1999 to 2016 (Data source: ITC, CHP, DH)**

Year	Health care workers		Non- Health care workers		Total	
	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)
Jul-Dec 1999	2	0 (0.0%)	3	0 (0.0%)	5	0 (0.0%)
2000	15	0 (0.0%)	20	1 (5.0%)	35	1 (2.9%)
2001	22	0 (0.0%)	50	1 (2.0%)	72	1 (1.4%)
2002	27	0 (0.0%)	50	1 (2.0%)	77	1 (1.3%)
2003	18	0 (0.0%)	43	0 (0.0%)	61	0 (0.0%)
2004	17	0 (0.0%)	40	0 (0.0%)	57	0 (0.0%)
2005	10	0 (0.0%)	57	0 (0.0%)	67	0 (0.0%)
2006	33	0 (0.0%)	139	0 (0.0%)	172	0 (0.0%)
2007	36	0 (0.0%)	118	0 (0.0%)	154	0 (0.0%)
2008	23	0 (0.0%)	126	3 (2.4%)	149	3 (2.0%)
2009	25	0 (0.0%)	161	1 (0.6%)	186	1 (0.5%)
2010	25	0 (0.0%)	131	0 (0.0%)	156	0 (0.0%)
2011	17	0 (0.0%)	145	0 (0.0%)	162	0 (0.0%)
2012	37	0 (0.0%)	154	0 (0.0%)	191	0 (0.0%)
2013	26	0 (0.0%)	162	1 (0.6%)	188	1 (0.5%)
2014	29	0 (0.0%)	157	0 (0.0%)	186	0 (0.0%)
2015	34	0 (0.0%)	150	0 (0.0%)	184	0 (0.0%)
2016	47	1 (2.1%)	145	1 (0.7%)	192	2 (1.0%)
Total	443	1 (0.2%)	1851	9 (0.5%)	2294	10 (0.4%)

## HEPATITIS C

**Box 51. Prevalence of anti-HCV in drug users on rehabilitation (Data source: PHLBSB, CHP, DH)**

Year	No. tested	Anti-HCV +ve (%)
1988/1989	134	99 (73.9%)
2000/2001	210	97 (46.2%)

**Box 52. Prevalence of anti-HCV at baseline screening of HIV/AIDS patients attending ITC from 2000 to 2017 (Data source: ITC, CHP, DH)**

Year	Male		Female		Total	
	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)
2000	54	5 (9.3%)	15	0 (0.0%)	69	5 (7.2%)
2001	72	9 (12.5%)	22	1 (4.5%)	94	10 (10.6%)
2002	118	9 (7.6%)	23	1 (4.3%)	141	10 (7.1%)
2003	89	13 (14.6%)	14	0 (0.0%)	103	13 (12.6%)
2004	108	21 (19.4%)	21	3 (14.3%)	129	24 (18.6%)
2005	137	19 (13.9%)	31	1 (3.2%)	168	20 (11.9%)
2006	187	49 (26.2%)	23	3 (13.0%)	210	52 (24.8%)
2007	215	41 (19.1%)	27	1 (3.7%)	242	42 (17.4%)
2008	201	40 (19.9%)	33	3 (9.1%)	234	43 (18.4%)
2009	168	33 (19.6%)	27	1 (3.7%)	195	34 (17.4%)
2010	163	15 (9.2%)	33	0 (0.0%)	196	15 (7.7%)
2011	168	12 (7.1%)	33	4 (12.1%)	201	16 (8.0%)
2012	226	10 (4.4%)	45	2 (4.4%)	271	12 (4.4%)
2013	264	11 (4.2%)	40	0 (0.0%)	304	11 (3.6%)
2014	301	5 (1.7%)	31	0 (0.0%)	332	5 (1.5%)
2015	327	14 (4.3%)	26	1 (3.8%)	353	15 (4.2%)
2016	299	21 (7.0%)	25	0 (0.0%)	324	21 (6.5%)
2017	331	16 (4.8%)	32	1 (3.1%)	363	17 (4.7%)

## HEPATITIS C

**Box 53. Prevalence of anti-HCV per HIV risk at baseline screening of HIV/AIDS patients attending ITC from 2000 to 2017 (Data source: ITC, CHP, DH)**

HIV risk	No. tested	Anti-HCV +ve (%)
Heterosexual male	792	50* (6.3%)
Heterosexual female	468	8 (1.7%)
Homo/Bi-sexual	2354	50 (2.1%)
Drug user	258	252 (97.7%)
Blood/blood product recipient	14	4 (28.6%)
Perinatal	9	0 (0.0%)
Undetermined	34	1 (2.9%)
Total	3929	365 (9.3%)

*\*30 out of 50 had a past history of injecting drug use*

## HEPATITIS C

**Box 54. Prevalence of anti-HCV from screening of blood donors and clinical testing of patients in 2 hospital clusters under Hospital Authority from 2007 to 2017 (Data source: HKRCBTS, PMH Microbiology Laboratory, PWH Microbiology Laboratory)**

Category	2007		2008		2009		2010		2011		2012		2013		2014		2015		2016		2017		Overall	
	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)	No. tested	Anti-HCV +ve (%)
1. BLOOD DONATION	205682	42 (< 0.1%)	211963	52 (< 0.1%)	231375	47 (< 0.1%)	226775	40 (< 0.1%)	234444	51 (< 0.1%)	243525	37 (< 0.1%)	247069	46 (< 0.1%)	254087	31 (< 0.1%)	260429	33 (< 0.1%)	257262	28 (< 0.1%)	243667	25 (< 0.1%)	2616278	432 (< 0.1%)
2. SCREENING																								
Pre-transplant	31	1 (3.2%)	18	0 (0.0%)	48	1 (2.1%)	68	2 (2.9%)	80	0 (0.0%)	96	0 (0.0%)	82	0 (0.0%)	111	1 (0.9%)	118	0 (0.0%)	108	0 (0.0%)	128	0 (0.0%)	888	5 (0.6%)
Drug users	118	29 (24.6%)	134	66 (49.3%)	154	93 (60.4%)	116	75 (64.7%)	84	61 (72.6%)	103	53 (51.5%)	112	63 (56.3%)	114	66 (57.9%)	124	51 (41.1%)	81	41 (50.6%)	87	38 (43.7%)	1227	636 (51.8%)
Needlestick injuries	546	6 (1.1%)	542	6 (1.1%)	574	5 (0.9%)	550	5 (0.9%)	559	4 (0.7%)	592	6 (1.0%)	610	4 (0.7%)	537	6 (1.1%)	494	3 (0.6%)	516	5 (1.0%)	667	9 (1.3%)	6187	59 (1.0%)
Haemodialysis/ peritoneal dialysis	1706	37 (2.2%)	1656	31 (1.9%)	1936	34 (1.8%)	2016	36 (1.8%)	2251	34 (1.5%)	2452	34 (1.4%)	2449	37 (1.5%)	2569	34 (1.3%)	2535	48 (1.9%)	2613	34 (1.3%)	3557	60 (1.7%)	25740	419 (1.6%)
Post-renal transplant	413	19 (4.6%)	470	21 (4.5%)	650	19 (2.9%)	680	25 (3.7%)	722	18 (2.5%)	737	17 (2.3%)	718	16 (2.2%)	692	15 (2.2%)	863	18 (2.1%)	541	6 (1.1%)	708	9 (1.3%)	7194	183 (2.5%)
Haematology (pre-chemotherapy)	223	0 (0.0%)	260	5 (1.9%)	262	2 (0.8%)	344	6 (1.7%)	399	1 (0.3%)	415	4 (1.0%)	444	2 (0.5%)	472	2 (0.4%)	489	4 (0.8%)	533	2 (0.4%)	687	6 (0.9%)	4528	34 (0.8%)
Rheumatology (pre-methotrexate)	210	1 (0.5%)	332	1 (0.3%)	396	5 (1.3%)	430	1 (0.2%)	464	2 (0.4%)	449	2 (0.4%)	471	4 (0.8%)	580	3 (0.5%)	689	5 (0.7%)	730	5 (0.7%)	1285	3 (0.2%)	6036	32 (0.5%)
History of blood transfusion	125	12 (9.6%)	197	18 (9.1%)	263	32 (12.2%)	239	21 (8.8%)	168	19 (11.3%)	197	17 (8.6%)	275	28 (10.2%)	224	22 (9.8%)	222	15 (6.8%)	166	14 (8.4%)	292	16 (5.5%)	2368	214 (9.0%)
Pre-vaccination	1	0 (0.0%)	1	0 (0.0%)	5	0 (0.0%)	0	0 (0.0%)	0	0 (0.0%)	0	0 (0.0%)	0	0 (0.0%)	0	0 (0.0%)	0	0 (0.0%)	0	0 (0.0%)	0	0 (0.0%)	7	0 (0.0%)
TOTAL (2)	3373	105 (3.1%)	3610	148 (4.1%)	4288	191 (4.5%)	4443	171 (3.8%)	4727	139 (2.9%)	5041	133 (2.6%)	5161	154 (3.0%)	5299	149 (2.8%)	5534	144 (2.6%)	5288	107 (2.0%)	7411	141 (1.9%)	54175	1582 (2.9%)
3. *CLINICAL INDICATION	4054	179 (4.4%)	5984	215 (3.6%)	7971	216 (2.7%)	8661	262 (3.0%)	8196	293 (3.6%)	9815	308 (3.1%)	10911	323 (3.0%)	11229	316 (2.8%)	12360	351 (2.8%)	15472	383 (2.5%)	15889	329 (2.1%)	110542	3175 (2.9%)
4. OTHERS OR UNKNOWN	8131	229 (2.8%)	8297	128 (1.5%)	7472	131 (1.8%)	8269	102 (1.2%)	8835	132 (1.5%)	9026	131 (1.5%)	9615	136 (1.4%)	11213	150 (1.3%)	10836	107 (1.0%)	10701	125 (1.2%)	15527	171 (1.1%)	107922	1542 (1.4%)
TOTAL (2+3+4)	15558	513 (3.0%)	17891	491 (2.7%)	19731	538 (2.7%)	21373	535 (2.5%)	21758	564 (2.6%)	23882	572 (2.4%)	25687	613 (2.4%)	27741	615 (2.2%)	28730	602 (2.1%)	31461	615 (2.0%)	38827	641 (1.7%)	272639	6299 (2.3%)

\*includes suspected hepatitis, work up for liver function derangement and others

## HEPATITIS C

**Box 55. Characteristics of anti-HCV positive subjects detected at HKRCBTS and 2 hospital clusters under Hospital Authority from 2005 to 2017 (Data source: HKRCBTS, PMH Microbiology Laboratory, PWH Microbiology Laboratory)**

		2005 (n=624)	2006 (n=542)	2007 (n=555)	2008 (n=543)	2009 (n=585)	2010 (n=575)	2011 (n=615)	2012 (n=609)	2013 (n=659)	2014 (n=646)	2015 (n=635)	2016 (n=643)	2017 (n=666)	Overall (n=7897)
		No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)
Lab	HKRCBTS	49 (7.9%)	35 (6.5%)	40 (7.2%)	49 (9.0%)	43 (7.4%)	38 (6.6%)	50 (6.6%)	35 (5.7%)	43 (6.5%)	31 (4.8%)	33 (5.2%)	28 (4.4%)	25 (3.8%)	499 (6.3%)
	PMH	229 (36.7%)	142 (26.2%)	89 (16.0%)	208 (38.3%)	273 (46.7%)	271 (47.1%)	280 (47.1%)	298 (48.9%)	279 (42.3%)	297 (46.0%)	354 (55.7%)	372 (57.9%)	340 (51.1%)	3432 (43.5%)
	PWH	346 (55.4%)	365 (67.3%)	426 (76.8%)	286 (52.7%)	269 (46.0%)	266 (46.3%)	285 (46.3%)	276 (45.3%)	337 (51.1%)	318 (49.2%)	248 (39.1%)	243 (37.8%)	301 (45.2%)	3966 (50.2%)
Sex	Male	413 (66.2%)	390 (72.0%)	377 (67.9%)	378 (69.6%)	415 (70.9%)	405 (70.4%)	434 (70.4%)	438 (71.9%)	464 (70.4%)	440 (68.1%)	434 (68.3%)	453 (70.5%)	454 (68.2%)	5495 (69.6%)
	Female	211 (33.8%)	152 (28.0%)	178 (32.1%)	165 (30.4%)	170 (29.1%)	170 (29.6%)	181 (29.6%)	171 (28.1%)	195 (29.6%)	206 (31.9%)	201 (31.7%)	190 (29.5%)	211 (31.7%)	2401 (30.4%)
	Unknown	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (0.2%)	1 (0.0%)
Age at diagnosis	Mean	46.8	47.4	50.3	49.8	52.9	51.2	50.8	51.1	51.0	52.0	54.0	54.6	55.7	51.4
	S.D.	15.9	16.6	16.3	17.9	16.9	17	16.5	16.3	16.6	16.2	15.5	15.5	15.1	16.3
	Range	0 - 87	0 - 101	0 - 94	0 - 88	1 - 102	0 - 90	0 - 90	0 - 99	0 - 113	0 - 95	1 - 95	0-97	0-94	0 - 113
Category	Blood donation	50 (8.0%)	35 (6.5%)	42 (7.6%)	52 (9.6%)	47 (8.0%)	40 (7.0%)	51 (8.3%)	37 (6.1%)	46 (7.0%)	31 (4.8%)	33 (5.2%)	28 (4.4%)	25 (3.8%)	517 (6.5%)
	Pre-transplant	2 (0.3%)	0 (0.0%)	1 (0.2%)	0 (0.0%)	1 (0.2%)	2 (0.3%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (0.2%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	7 (0.1%)
	Drug users	144 (23.1%)	59 (10.9%)	29 (5.2%)	66 (12.2%)	93 (15.9%)	75 (13.0%)	61 (9.9%)	53 (8.7%)	63 (9.6%)	66 (10.2%)	51 (8.0%)	41 (6.4%)	38 (5.7%)	839 (10.6%)
	Needlestick injuries	8 (1.3%)	7 (1.3%)	6 (1.1%)	6 (1.1%)	5 (0.9%)	5 (0.9%)	4 (0.7%)	6 (1.0%)	4 (0.6%)	6 (0.9%)	3 (0.5%)	5 (0.8%)	9 (1.4%)	74 (0.9%)
	Pre-haemodialysis/ peritoneal dialysis	40 (6.4%)	35 (6.5%)	37 (6.7%)	31 (5.7%)	34 (5.8%)	36 (6.3%)	34 (5.5%)	34 (5.6%)	37 (5.6%)	34 (5.3%)	48 (7.6%)	34 (5.3%)	60 (9.0%)	494 (6.3%)
	Post-renal transplant	17 (2.7%)	18 (3.3%)	19 (3.4%)	21 (3.9%)	19 (3.2%)	25 (4.3%)	18 (2.9%)	17 (2.8%)	16 (2.4%)	15 (2.3%)	18 (2.8%)	6 (0.9%)	9 (1.4%)	218 (2.8%)
	Haematology	3 (0.5%)	1 (0.2%)	0 (0.0%)	5 (0.9%)	2 (0.3%)	6 (1.0%)	1 (0.2%)	4 (0.7%)	2 (0.3%)	2 (0.3%)	4 (0.6%)	2 (0.3%)	6 (0.9%)	38 (0.5%)
	Pre-methotrexate	1 (0.2%)	1 (0.2%)	1 (0.2%)	1 (0.2%)	5 (0.9%)	1 (0.2%)	2 (0.3%)	2 (0.3%)	4 (0.6%)	3 (0.5%)	5 (0.8%)	5 (0.8%)	3 (0.5%)	34 (0.4%)
	History of blood transfusion	12 (1.9%)	11 (2.0%)	12 (2.2%)	18 (3.3%)	32 (5.5%)	21 (3.7%)	19 (3.1%)	17 (2.8%)	28 (4.2%)	22 (3.4%)	15 (2.4%)	14 (2.2%)	16 (2.4%)	237 (3.0%)
	Clinical Indication	155 (24.8%)	170 (31.4%)	179 (32.3%)	215 (39.6%)	216 (36.9%)	262 (45.6%)	293 (47.6%)	308 (50.6%)	323 (49.0%)	316 (48.9%)	351 (55.3%)	383 (59.6%)	329 (49.4%)	3500 (44.3%)
	Others or unknown	192 (30.8%)	205 (37.8%)	229 (41.3%)	128 (23.6%)	131 (22.4%)	102 (17.7%)	132 (21.5%)	131 (21.5%)	136 (20.6%)	150 (23.2%)	107 (16.9%)	125 (19.4%)	171 (25.7%)	1939 (24.6%)

# SURVEILLANCE OF VIRAL HEPATITIS IN HONG KONG

## 2017 UPDATE

### Liver cancers


#### Tabulated results

<b>Box</b>	<b>Title</b>	<b>Page</b>
Box 56.	Hong Kong liver cancer statistics, number of new cases by gender from 2001 - 2016 (Data source: Hong Kong Cancer Registry, Hospital Authority)	81
Box 57.	Hong Kong liver cancer statistics, number of new cases and incidence rate by age and gender, from 2001 – 2016 (Data source: Hong Kong Cancer Registry, Hospital Authority)	82
Box 58.	Hong Kong liver cancer mortality statistics by gender from 2001 - 2016 (Data source: Hong Kong Cancer Registry, Hospital Authority)	83
Box 59.	Hong Kong liver cancer mortality statistics, by age and gender, from 2001 – 2016 (Data source: Hong Kong Cancer Registry, Hospital Authority)	84



## Liver Cancer

Box 56. Hong Kong liver cancer statistics, number of new cases by gender from 2001 - 2016 (Data source: Hong Kong Cancer Registry, Hospital Authority)



	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Female	324	343	390	385	425	414	386	426	447	465	459	426	445	478	435	419
Male	1313	1233	1264	1278	1324	1331	1304	1319	1385	1398	1399	1364	1407	1369	1356	1391
Total	1637	1576	1654	1663	1749	1745	1690	1745	1832	1863	1858	1790	1852	1847	1791	1810

## Liver Cancer

**Box 57. Hong Kong liver cancer statistics, number of new cases and incidence rate by age and gender, from 2001 – 2016 (Data source: Hong Kong Cancer Registry, Hospital Authority)**

Year	0-19						20-44						45-64						65+						Crude rate			ASR		
	Male		Female		Total		Male		Female		Total		Male		Female		Total		Male		Female		Total		Male	Female	Total	Male	Female	Total
	N	I	N	I	N	I	N	I	N	I	N	I	N	I	N	I	N	I	N	I	N	I	N	I	CR	CR	CR	ASR	ASR	ASR
2001	4	0.5	1	0.1	5	0.3	130	9.5	26	1.7	156	5.3	590	76.9	86	12.1	676	45.7	589	169.3	211	52.0	800	106.2	40.0	9.4	24.4	32.7	7.4	20.1
2002	4	0.5	2	0.3	6	0.4	130	9.7	17	1.1	147	5.1	534	67.1	79	10.5	613	39.5	565	157.6	245	58.5	810	104.2	37.6	9.9	23.4	30.0	7.4	18.6
2003	6	0.8	2	0.3	8	0.5	110	8.4	25	1.6	135	4.7	581	70.5	100	12.6	681	42.1	567	154.5	263	61.4	830	104.4	38.8	11.2	24.6	30.3	8.2	19.1
2004	2	0.3	1	0.1	3	0.2	121	9.4	18	1.2	139	4.9	554	64.6	91	10.9	645	38.1	601	159.2	275	62.3	876	107	39.1	10.9	24.5	29.6	7.8	18.4
2005	2	0.3	0	0	2	0.1	110	8.7	21	1.4	131	4.7	605	67.5	110	12.4	715	40.1	607	157.8	294	65.3	901	107.9	40.6	12.0	25.7	29.9	8.3	18.9
2006	6	0.8	1	0.1	7	0.5	88	7.1	21	1.4	109	3.9	637	68.5	109	11.8	746	40.2	600	152.6	283	61.7	883	103.6	40.7	11.5	25.4	29.3	8.0	18.4
2007	2	0.3	1	0.2	3	0.2	83	6.8	13	0.8	96	3.5	621	64.7	95	9.8	716	37.1	598	148.3	277	59.1	875	100.3	39.7	10.6	24.4	27.9	7.1	17.2
2008	1	0.1	1	0.2	2	0.1	90	7.5	24	1.6	114	4.2	636	64	135	13.2	771	38.3	592	144.6	266	56.2	858	97.2	40.1	11.6	25.1	27.4	7.6	17.2
2009	2	0.3	2	0.3	4	0.3	87	7.4	20	1.3	107	4	695	68	131	12.3	826	39.6	601	143.8	294	61.1	895	99.6	42.2	12.1	26.3	27.9	7.7	17.5
2010	0	0	4	0.7	4	0.3	78	6.7	23	1.5	101	3.8	711	67.9	140	12.6	851	39.5	609	142.4	298	60.7	907	98.7	42.4	12.5	26.5	27.1	8.1	17.3
2011	6	0.9	3	0.5	9	0.7	85	7.4	22	1.5	107	4	694	65	122	10.7	816	36.9	614	140.1	312	62.0	926	98.4	42.4	12.2	26.3	26.8	7.5	16.8
2012	2	0.3	1	0.2	3	0.2	69	6.0	25	1.6	94	3.5	654	60.6	108	9.2	762	33.9	639	140.1	292	55.7	931	95.0	41.0	11.1	25.0	25.1	6.5	15.5
2013	6	1	2	0.3	8	0.7	64	5.6	19	1.2	83	3.1	698	64.3	126	10.6	824	36.2	639	134.5	298	54.7	937	91.9	42.2	11.6	25.8	25.4	6.9	15.8
2014	3	0.5	1	0.2	4	0.3	69	6	17	1.1	86	3.2	644	59.2	130	10.8	774	33.7	653	131.7	330	58.1	983	92.4	40.9	12.3	25.5	23.8	6.9	15
2015	1	0.2	2	0.3	3	0.3	51	4.4	14	0.9	65	2.4	621	57.2	107	8.7	728	31.5	683	131.3	312	52.5	995	89.3	40.3	11.1	24.6	22.7	6.2	14.1
2016	1	0.2	2	0.4	3	0.3	64	5.6	9	0.6	73	2.7	679	62.6	118	9.5	797	34.2	647	119.2	290	46.8	937	80.6	41.2	10.6	24.7	23	5.7	13.9
Average	3	0.4	2	0.3	5	0.3	89	7.3	20	1.3	109	4.0	635	65.2	112	11.0	746	37.5	613	143.9	284	57.7	897	97.7	40.6	11.3	25.1	27.2	7.2	16.9

Notes:

I: Incidence rate per 100000 population


N: No. of new cases by selected age groups

ASR: Age-standardized rate (per 100000 population) is calculated based on the reference standard population used

CR: Crude rate per 100000 population

## Liver Cancer

**Box 58. Hong Kong liver cancer mortality statistics by gender from 2001 - 2016 (Data source: Hong Kong Cancer Registry, Hospital Authority)**



	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Female	353	291	337	341	387	387	351	383	416	417	396	460	401	436	432	405
Male	1071	1090	1075	1076	1119	1075	1098	1116	1072	1113	1140	1045	1123	1149	1139	1135
Total	1424	1381	1412	1417	1506	1462	1449	1499	1488	1530	1536	1505	1524	1585	1571	1540

## Liver Cancer

**Box 59. Hong Kong liver cancer mortality statistics, by age and gender, from 2001 – 2016 (Data source: Hong Kong Cancer Registry, Hospital Authority)**

Year	0-19						20-44						45-64						65+						Crude rate			ASR		
	Male		Female		Total		Male		Female		Total		Male		Female		Total		Male		Female		Total		Male	Female	Total	Male	Female	Total
	N	I	N	I	N	I	N	I	N	I	N	I	N	I	N	I	N	I	N	I	N	I	N	I	CR	CR	CR	ASR	ASR	ASR
2001	3	0.4	2	0.3	5	0.3	101	7.4	16	1	117	4	434	56.6	74	10.4	508	34.3	533	153.2	261	64.4	794	105.4	32.6	10.3	21.2	26.8	7.8	17.1
2002	3	0.4	1	0.1	4	0.3	98	7.3	15	1	113	3.9	425	53.4	51	6.7	476	30.7	564	157.3	224	53.5	788	101.4	33.2	8.4	20.5	26.4	5.9	16.1
2003	2	0.3	0	0	2	0.1	80	6.1	15	1	95	3.3	436	52.9	69	8.7	505	31.2	557	151.8	253	59	810	101.8	33	9.7	21	25.6	6.8	15.9
2004	2	0.3	0	0	2	0.1	66	5.1	15	1	81	2.9	428	49.9	69	8.2	497	29.3	580	153.6	257	58.2	837	102.2	32.9	9.7	20.9	24.7	6.6	15.4
2005	0	0	1	0.1	1	0.1	93	7.4	17	1.1	110	3.9	432	48.2	75	8.5	507	28.5	594	154.4	294	65.3	888	106.4	34.3	10.9	22.1	24.8	7.2	15.8
2006	2	0.3	0	0	2	0.1	49	3.9	12	0.8	61	2.2	420	45.2	64	6.9	484	26.1	604	153.6	311	67.8	915	107.4	32.9	10.8	21.3	23.3	6.7	14.7
2007	3	0.4	0	0	3	0.2	57	4.7	7	0.5	64	2.3	470	49	62	6.4	532	27.6	568	140.8	282	60.1	850	97.5	33.4	9.7	21	23.1	5.9	14.2
2008	1	0.1	0	0	1	0.1	68	5.7	17	1.1	85	3.1	480	48.3	82	8	562	27.9	567	138.5	284	60	851	96.4	33.9	10.4	21.5	22.9	6.3	14.3
2009	2	0.3	0	0	2	0.2	43	3.7	10	0.7	53	2	442	43.3	95	8.9	537	25.7	585	140	311	64.7	896	99.7	32.6	11.3	21.3	21.2	6.7	13.7
2010	0	0	0	0	0	0	35	3	15	1	50	1.9	474	45.3	89	8	563	26.1	604	141.2	313	63.8	917	99.8	33.8	11.2	21.8	21.2	6.5	13.6
2011	1	0.2	1	0.2	2	0.2	52	4.5	8	0.5	60	2.2	462	43.3	72	6.3	534	24.1	625	142.6	315	62.6	940	99.9	34.5	10.5	21.7	21.2	5.9	13.2
2012	0	0	1	0.2	1	0.1	50	4.3	10	0.7	60	2.2	431	39.9	95	8.1	526	23.4	564	123.7	354	67.6	918	93.7	31.4	12	21	18.9	6.5	12.4
2013	3	0.5	1	0.2	4	0.3	38	3.3	13	0.8	51	1.9	437	40.2	82	6.9	519	22.8	645	135.8	305	56.0	950	93.1	33.7	10.4	21.2	19.4	5.6	12.1
2014	2	0.3	0	0	2	0.2	48	4.2	11	0.7	59	2.2	469	43.1	71	5.9	540	23.5	629	126.8	354	62.3	983	92.4	34.3	11.2	21.9	19.4	5.7	12.2
2015	1	0.2	1	0.2	2	0.2	37	3.2	6	0.4	43	1.6	427	39.4	76	6.2	503	21.8	674	129.6	349	58.7	1023	91.8	33.8	11.0	21.5	18.4	5.4	11.6
2016	1	0.2	1	0.2	2	0.2	39	3.4	7	0.5	46	1.7	445	41.1	75	6.0	520	22.3	650	119.7	322	51.9	972	83.6	33.6	10.2	21.0	18.0	4.9	11.0
Average	2	0.2	<1	0.1	2	0.2	60	4.9	12	0.8	72	2.6	445	45.6	75	7.4	520	26.1	596	140.0	299	60.8	896	97.6	33.4	10.5	21.3	22.0	6.2	13.8

Notes:

I: Mortality rate per 100000 population

N: No. of death cases by selected age groups

ASR: Age-standardized rate (per 100000 population) is calculated based on the reference standard population used

CR: Crude rate per 100000 population

# SURVEILLANCE OF VIRAL HEPATITIS IN HONG KONG

## 2017 UPDATE

### ABBREVIATIONS

AIDS	Acquired immune deficiency syndrome
Anti-HAV	Antibody against hepatitis A virus
Anti-HBc	Antibody against hepatitis B core antigen
Anti-HBs	Antibody against hepatitis B surface antigen
Anti-HCV	Antibody against hepatitis C virus
Anti-HEV	Antibody against hepatitis E virus
BUHC	Baptist University Health Centre
CHP	Centre for Health Protection
CRPVH	Community Research Project on Viral Hepatitis
CUHC	City University Health Centre
CUHK	Chinese University of Hong Kong
DH	Department of Health
FHS	Family Health Service
FPA	Family Planning Association
HBsAg	Hepatitis B surface antigen
HAV	Hepatitis A virus
HBV	Hepatitis B virus
HCC	Hepatocellular carcinoma
HCV	Hepatitis C virus
HCW	Health care worker
HEV	Hepatitis E virus
HIV	Human immunodeficiency virus
HKRCBTS	Hong Kong Red Cross Blood Transfusion Service
IgM	Immunoglobulin M
IDU	Injecting drug users
ITC	Integrated Treatment Centre
LUHC	Lingnan University Health Centre
MCHC	Maternal and Child Health Centre
MSM	Men who have sex with men
PHIS	Public Health Information System
PHLSB	Public Health Laboratory Services Branch
PMH	Princess Margaret Hospital
PWH	Prince of Wales Hospital
SEB	Surveillance and Epidemiology Branch
TPC	Therapeutic Prevention Clinic

# SURVEILLANCE OF VIRAL HEPATITIS IN HONG KONG

## 2017 UPDATE

### REFERENCES

1. Gust ID. The epidemiology of viral hepatitis. In: Vyas GN, Dienstag JL, Hoofnagle JH, editors: *Viral Hepatitis and Liver Disease*. Orlando: Grune & Stratton;1984. p. 415-21.
2. Wong KH, Liu YM, Ng PS, et al. Epidemiology of hepatitis A and hepatitis E infection and their determinants in adult Chinese community in Hong Kong. *J Med Virol* 2004;72:538-44.
3. Poon C, Ho B. Update of Hepatitis A in Hong Kong. *CD Watch* 2015;12:14.
4. Centre for Health Protection, Department of Health. Summary on the recent upsurge of hepatitis A infection among MSM in Hong Kong. *CD Watch* 2017;14:24.
5. Chin KP, Lok ASF, Wong LSK, et al. Current seroepidemiology of hepatitis A in Hong Kong. *J Med Virol* 1991;34:191-3.
6. Tsang CW, Chan CL. Epidemiology of viral hepatitis in Hong Kong. In: *New trends in peptic ulcer and chronic hepatitis-Part II. Chronic Hepatitis*. Tokyo: Excerpta Medica;1987. p. 43-50.
7. Centre for Health Protection, Department of Health. Seroprevalence rates of hepatitis A virus antibodies. (Available at <https://www.chp.gov.hk/en/statistics/data/10/641/701/3936.html>. Accessed 14 September 2017)
8. Scientific Committee on AIDS and STI and Scientific Committee on Vaccine Preventable Diseases. Statement on hepatitis A vaccination and men who have sex with men. June 2017 (Available at [https://www.chp.gov.hk/files/pdf/statement\\_on\\_hav\\_and\\_msm\\_201706.pdf](https://www.chp.gov.hk/files/pdf/statement_on_hav_and_msm_201706.pdf). Accessed 30 October 2018)
9. Chiu DM, Chan MC, Yeung AC. Seroprevalence of hepatitis E virus in Hong Kong, 2008-2009. *J Med Virol*. 2013;85(3):459-61.
10. Centre for Health Protection, Department of Health. Review of hepatitis E infection (2001-2010). *CD Watch* 2011;8:1.
11. Centre for Health Protection, Department of Health. Update of hepatitis E infection in Hong Kong. *CD Watch* 2017;14:13.
12. Chau TN, Lai ST, Tse C, et al. Epidemiology and clinical features of sporadic hepatitis E as compared with hepatitis A. *Am J Gastroenterol* 2006;101:292-6.
13. Lam WY, Chan RCW, Sung JJY, et al. Genotype distribution and sequence variation of hepatitis E virus, Hong Kong. *Emerging Infectious Diseases* 2009;15:792-4.

14. Centre for Health Protection, Department of Health. Update in hepatitis E infection in Hong Kong. *CD Watch* 2012;9:5.
15. Chow CW, Tsang SW, Tsang OT, et al. Comparison of acute hepatitis E infection outcome in patients with and without chronic hepatitis B infection: a 10 year retrospective study in three regional hospitals in Hong Kong. *J Clin Virol*. 2014;60(1):4-10.
16. Centre for Food Safety, Food and Environmental Hygiene Department. Hepatitis E Virus in Fresh Pig Livers. *Risk Assessment Studies Report HKSAR* 2010;44:39.
17. Hepatitis E vaccine: WHO position paper, May 2015. Available at <http://www.who.int/wer/2015/wer9018.pdf>. Accessed on 14 September 2015.
18. Kwan LC, Ho YY, Lee SS. The declining HBsAg carriage rate in pregnant women in Hong Kong. *Epidemiol Infect* 1997;119:281-3.
19. Cooley L, Sasadeusz J. Clinical and virological aspects of hepatitis B co-infection in individuals infected with human immunodeficiency virus type-1. *J Clin Virol* 2003;26:185-93.
20. Yuen MF, Sablon E, Tanaka Y, et al. Epidemiological study of hepatitis B virus genotypes, core promoter and precore mutations of chronic hepatitis B infection in Hong Kong. *J Hepatol* 2004;41:119-25.
21. Chan HL, Hui AY, Wong ML, et al. Genotype C hepatitis B virus infection is associated with an increased risk of hepatocellular carcinoma. *Gut* 2004;53:1494-8.
22. Chan HL, Tsui SK, Tse CH, et al. Epidemiological and virological characteristics of 2 subgroups of hepatitis B virus genotype C. *J Infect Dis* 2005;191:2022-32.
23. Zhu L, Tse CH, Wong VSW, et al. A complete genomic analysis of hepatitis B virus genotypes and mutations in HbeAg-negative chronic hepatitis B in China. *J Viral Hepatol* 2008;15:449-58.
24. Chan HL, Tse CH, Mo G, et al. High viral load and hepatitis B virus subgenotype Ce are associated with increased risk of hepatocellular carcinoma. *J Clin Oncol* 2008;26:177-82.
25. Chan HL, Wong GL, Tse CH et al. Hepatitis B virus genotype C is associated with more severe liver fibrosis than genotype B. *Clin Gastroenterol Hepatol* 2009;7:1361-6.
26. Wong GL, Chan HL, Yiu KK, et al. Meta-analysis: The association of hepatitis B virus genotypes and hepatocellular carcinoma. *Aliment Pharmacol Ther*. 2013;37:517-26.
27. Lo CM, Cheung CK, Lau GK, et al. Significance of hepatitis B virus genotype in liver transplantation for chronic hepatitis B. *Am J Transplant* 2005;5:1893-900.
28. Yuen MF, Tanaka Y, Mizokami M, et al. Role of hepatitis B virus genotypes Ba and C, core promoter and precore mutations on hepatocellular carcinoma: a case control study. *Carcinogenesis* 2004;25:1593-8.
29. Chan AO, Yuen MF, Lam CM, et al. Prevalence and characteristics of familial hepatocellular carcinoma caused by chronic hepatitis B infection in Hong Kong. *Aliment Pharmacol Ther* 2004;19:401-6.

30. Ho CF, Wong KH, Chan CW, et al. Current pattern and course of acute hepatitis B virus infection in Hong Kong. *J Gastroenterol Hepatol* 2003;19:602-3.
31. Young BWY, Lee SS, Lim WL, et al. The long-term efficacy of plasma-derived hepatitis B vaccine in babies born to carrier mothers. *J Viral Hepatol* 2003;10:23-30.
32. Lin AWC, Wong KH. Long-term protection of neonatal hepatitis B vaccination in a 30-year cohort in Hong Kong. *J Hepatol* 2013;59:1363-4.
33. Yuen MF, Lim WL, Chan AO, et al. 18-year follow-up study of a prospective randomized trial of hepatitis B vaccinations without booster doses in children. *Clin Gastroenterol Hepatol* 2004;2:941-5.
34. Tse W, Mok T. Survey on Immunisation coverage among children aged two to five. *Public Health & Epidemiology Bulletin* 2002;11:13-8.
35. Tse WKM, Yeung SWT. Immunisation coverage among children aged two to five: an update. *Public Health & Epidemiology Bulletin* 2004;13:7-15.
36. Wu T, Chan SK, Kung KH, et al. Immunization coverage among children aged two to five: findings of the 2006 survey. *Public Health & Epidemiology Bulletin* 2007;16:57-68.
37. Chung PW, Suen SH, Chan OK, et al. Awareness and knowledge of hepatitis B infection and prevention and the use of hepatitis B vaccination in the Hong Kong adult Chinese population. *Chin Med J* 2012;125:422-7.
38. Centre for Health Protection, Department of Health. Hepatitis C in Hong Kong, 2008 to 2011. *CD Watch* 2011;8:25.
39. Hagan H, Jordan AE, Neurer J, et al., Incidence of sexually transmitted hepatitis C virus infection in HIV-positive men who have sex with men. *AIDS* 2015 Nov;29(17):2335-45.
40. Chan GCB, Lim WL, Yeoh EK. Prevalence of hepatitis C infection in Hong Kong. *J Gastroen Hepatol* 1992;7:117-20.
41. Wong HK, Lee CK, Leung JN, et al. Risk factor analysis of hepatitis C virus infection among Chinese blood donors in Hong Kong. *Transfus Med* 2012 Apr;22(2):133-6.
42. Au WY, Lee V, Kho B, et al. A synopsis of current haemophilia care in Hong Kong. *Hong Kong Med J* 2011 Jun;17(3):189-94.
43. Chan TM, Lok AS, Cheng IK, et al. Prevalence of hepatitis C virus infection in hemodialysis patients: a longitudinal study comparing the results of RNA and antibody assays. *Hepatology* 1993;17:5-8.
44. Lee KCK, Lim WWL, Lee SS. High prevalence of HCV in a cohort of injectors on methadone substitution treatment. *J Clin Virol* 2008;41:297-300.
45. Wong NS, Chan PC, Lee SS, et al., A multilevel approach for assessing the variability of hepatitis C prevalence in injection drug users by their gathering places. *Int J Infect Dis* 2013 Mar;17(3):e193-8.



46. Wong VW, Wong GL, Chim AM, et al., Targeted hepatitis C screening among ex-injection drug users in the community *J Gastroenterol Hepatol*. 2014 Jan; 29(1): 116-20.
47. Centers for Disease Control and Prevention (CDC). Sexual Transmission of Hepatitis C Virus Among HIV-Infected Men Who Have Sex with Men --- New York City, 2005--2010. *MMWR Morb Mortal Wkly Rep*. 2011 Jul 22;60:945-50.
48. Lin A, Wong P, Lo J. A case series of hepatitis C infection and syphilis among HIV positive men who have sex with men. *Communicable Disease Watch* 2014; 11.
49. Lin AWC, Wong KH, Chan K, More safer sex intervention needed for HIV-positive MSM with higher education level for prevention of sexually transmitted hepatitis C. Poster presentation P 131; HIV Drug Therapy Glasgow 2014.
50. Monga HK, Rodriguez-Barradas MC, Breaux K, et al. Hepatitis C virus infection-related morbidity and mortality among patients with human immunodeficiency virus infection. *Clin Infect Dis* 2001;33:240-7.
51. Delwart E, Slikas E, Stramer SL, et al. Genetic Diversity of Recently Acquired and Prevalent HIV, Hepatitis B Virus and Hepatitis C Virus Infections in US Blood Donors. *JID* 2012; 205:875-85.
52. Prescott LE, Simmonds P, Lai CL, et al. Detection and clinical features of hepatitis C virus type 6 infections in blood donors from Hong Kong. *J Med Virol* 1996;50:168-75.
53. Wong DA, Tong LK, Lim W. High prevalence of hepatitis C virus genotype 6 among certain risk groups in Hong Kong. *Eur J Epidemiol* 1998;14:421-6.
54. Chan TM, Lau JYN, Wu PC, et al. Hepatitis C virus genotypes in patients on renal replacement therapy. *Nephrol Dial Transplant* 1998;13:731-4.
55. Zhou DX, Tang JW, Chu IM, et al. Hepatitis C virus genotype distribution among intravenous drug user and the general population in Hong Kong. *J Med Virol* 2006;78:574-81.
56. Chan D, Lee SS, Lee KC. The effects of widespread methadone treatment on the molecular epidemiology of hepatitis C virus infection among injecting drug users in Hong Kong. *J Med Virol* 2011;83:1187-94.

57. Chan DP, Lin AW, Wong KH, et al., Diverse origins of hepatitis C virus in HIV co-infected men who have sex with men in Hong Kong. *Virology* 2015; 12:120
58. Seto WK, Lai CL, Fung J, et al. Natural history of chronic hepatitis C: genotype 1 versus genotype 6. *J Hepatol* 2010;53:444-8.
59. World Health Organization. Cancer fact sheet. (Available at <http://www.who.int/mediacentre/factsheets/fs297/en/>. Accessed on 2 September 2017)
60. Yuen MF, Hou JL, Chutaputti A, et al. Hepatocellular carcinoma in the Asia Pacific Region. *J Gastroenterol Hepatol* 2009;24:346-353.
61. Lo CM, Fan ST, Liu CL, et al. Ten-year experience with liver transplantation at Queen Mary Hospital: retrospective study. *Hong Kong Med J* 2002;8:240-4.
62. Hong Kong Cancer Registry, Hospital Authority. (Available at <http://www3.ha.org.hk/cancereg/pdf/overview/Summary%20of%20CanStat%202016.pdf>. Accessed on 28 December 2018)
63. Lim WL, Yeoh EK. Hepatitis A vaccination. *Lancet* 1992;339:304.
64. Lai CL. Hepatitis A risk heightened. Data quoted in *United Daily News* dated 10 June 1994.
65. Lee A, Cheng F, Lau L, et al. Changing hepatitis A epidemiology among Hong Kong Chinese adolescents: what are the implications? *Public Health* 1999; 113:185-8.